

HIV SEMI-ANNUAL SURVEILLANCE REPORT

San Francisco Department of Public Health

HIV Cases Reported Through June 2015

Contents	Page
Surveillance Summary.....	1
Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	2
Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	2
Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis.....	3
Table 2: Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis.....	4
Table 3: Diagnoses of HIV infection by gender and year of initial HIV diagnosis	5
Table 4: Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis...	6
Table 5: Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis.....	7
Table 6: Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis.....	8
Table 7: Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis.....	9
Table 8: Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis.....	10
Table 9: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	11
Table 10: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	12
Table 11: Persons living with HIV infection by age, gender and race/ethnicity.....	13
Table 12: Persons living with HIV infection by transmission category, gender and race/ethnicity.....	14

The HIV Surveillance Report is published semi-annually by the
San Francisco Department of Public Health, Population Health Division
Applied Research, Community Health Epidemiology, and Surveillance Branch (ARCHES)
25 Van Ness Avenue, Suite 500, San Francisco, CA 94102; Phone (415) 437-6336, FAX (415) 431-0353

Director of Health: Barbara A. Garcia, MPA
Director of Population Health Division: Tomas Aragon, MD, DrPH
HIV Surveillance Section Directors: Susan Scheer, PhD, MPH; Ling Hsu, MPH

***The HIV Surveillance Report is accessible via internet:
<http://www.sfdph.org/dph/files/reports/default.asp>***

HIV SURVEILLANCE SUMMARY

San Francisco¹ (as of 06/30/2015)

Persons living with HIV infection ²	16,014
Diagnoses of HIV infection stage 3 (AIDS), cumulative	29,741
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	20,196

California³ (as of 12/31/2013)

Persons living with HIV infection ²	119,878
Diagnoses of HIV infection stage 3 (AIDS), cumulative	169,734
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	96,443

United States⁴ (as of 12/31/2012)

Persons living with HIV infection ²	929,646
Diagnoses of HIV infection stage 3 (AIDS), cumulative (as of 12/31/2013)	1,217,863
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	676,361


¹ Residents of San Francisco at time of HIV or AIDS diagnosis.

² Persons diagnosed with HIV infection regardless of disease stage.

³ CA data source: www.cdph.ca.gov/data/statistics/Pages/OAHIVAIDSStatistics.aspx.


⁴ US data source: CDC HIV Surveillance Report 2013, vol. 25 (Table 2b, 11b, 14b).
http://www.cdc.gov/hiv/library/reports/surveillance/2013/surveillance_Report_vol_25.html.

Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2015*



* Reporting for recent years is incomplete. See Table 9 for actual numbers per year.

Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2015 †



† Reporting for recent years is incomplete. See Table 10 for actual numbers per year.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2015

Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis (1), San Francisco, 2004-2015

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)
0 - 12	2 (0.3)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.1)	1 (0.2)	1 (0.2)	2 (0.4)	5 (1.0)	1 (0.2)
18 - 24	62 (8.8)	52 (9.5)	59 (11.4)	55 (10.4)	53 (10.2)	56 (12.0)
25 - 29	92 (13.0)	70 (12.7)	68 (13.1)	100 (18.9)	83 (16.0)	58 (12.5)
30 - 39	282 (39.9)	208 (37.8)	175 (33.7)	190 (35.9)	183 (35.2)	144 (31.0)
40 - 49	189 (26.8)	155 (28.2)	143 (27.6)	127 (24.0)	149 (28.7)	127 (27.3)
50 - 59	61 (8.6)	51 (9.3)	56 (10.8)	42 (7.9)	40 (7.7)	62 (13.3)
60 +	17 (2.4)	12 (2.2)	17 (3.3)	13 (2.5)	7 (1.3)	17 (3.7)
Total	706 (100)	550 (100)	519 (100)	529 (100)	520 (100)	465 (100)

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	2 (0.5)	1 (0.2)	0 (0.0)	0 (0.0)	1 (0.3)	0 (0.0)
18 - 24	55 (12.5)	44 (10.7)	53 (12.4)	50 (13.4)	36 (11.8)	18 (16.2)
25 - 29	56 (12.7)	64 (15.5)	71 (16.6)	80 (21.4)	50 (16.3)	24 (21.6)
30 - 39	137 (31.1)	107 (26.0)	130 (30.4)	110 (29.4)	94 (30.7)	24 (21.6)
40 - 49	125 (28.3)	127 (30.8)	124 (29.0)	90 (24.1)	73 (23.9)	32 (28.8)
50 - 59	50 (11.3)	54 (13.1)	38 (8.9)	31 (8.3)	39 (12.7)	9 (8.1)
60 +	16 (3.6)	15 (3.6)	11 (2.6)	13 (3.5)	13 (4.2)	4 (3.6)
Total	441 (100)	412 (100)	427 (100)	374 (100)	306 (100)	111 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2015

Table 2. Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis (1), San Francisco, 2004-2015

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.2)	0 (0.0)	0 (0.0)	1 (0.2)	1 (0.2)	1 (0.3)
18 - 24	11 (2.3)	14 (2.9)	13 (2.9)	15 (3.4)	17 (3.9)	15 (4.6)
25 - 29	21 (4.3)	32 (6.6)	25 (5.6)	39 (8.7)	31 (7.1)	23 (7.1)
30 - 39	170 (35.0)	159 (33.0)	125 (27.9)	133 (29.8)	125 (28.5)	75 (23.1)
40 - 49	166 (34.2)	184 (38.2)	183 (40.8)	165 (37.0)	168 (38.4)	120 (36.9)
50 - 59	88 (18.1)	79 (16.4)	81 (18.1)	66 (14.8)	75 (17.1)	70 (21.5)
60 +	29 (6.0)	14 (2.9)	21 (4.7)	27 (6.1)	21 (4.8)	21 (6.5)
Total	486 (100)	482 (100)	448 (100)	446 (100)	438 (100)	325 (100)

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
18 - 24	15 (5.0)	12 (4.8)	10 (4.2)	9 (4.9)	4 (3.1)	3 (6.8)
25 - 29	27 (9.0)	20 (8.0)	25 (10.4)	17 (9.2)	13 (9.9)	2 (4.5)
30 - 39	73 (24.4)	62 (24.7)	60 (25.0)	42 (22.7)	40 (30.5)	6 (13.6)
40 - 49	99 (33.1)	98 (39.0)	80 (33.3)	58 (31.4)	37 (28.2)	15 (34.1)
50 - 59	62 (20.7)	50 (19.9)	46 (19.2)	39 (21.1)	24 (18.3)	11 (25.0)
60 +	23 (7.7)	9 (3.6)	19 (7.9)	20 (10.8)	13 (9.9)	7 (15.9)
Total	299 (100)	251 (100)	240 (100)	185 (100)	131 (100)	44 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2015

Table 3. Diagnoses of HIV infection by gender and year of initial HIV diagnosis (1), San Francisco, 2004-2015

Gender	Year of initial HIV diagnosis					
	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)
Male	638 (90.4)	489 (88.9)	469 (90.4)	462 (87.3)	462 (88.8)	422 (90.8)
Female	53 (7.5)	45 (8.2)	36 (6.9)	44 (8.3)	41 (7.9)	25 (5.4)
Transgender	15 (2.1)	16 (2.9)	14 (2.7)	23 (4.3)	17 (3.3)	18 (3.9)
Total	706 (100)	550 (100)	519 (100)	529 (100)	520 (100)	465 (100)

Gender	Year of initial HIV diagnosis					
	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)
Male	394 (89.3)	361 (87.6)	401 (93.9)	342 (91.4)	288 (94.1)	100 (90.1)
Female	34 (7.7)	41 (10.0)	22 (5.2)	22 (5.9)	14 (4.6)	9 (8.1)
Transgender	13 (2.9)	10 (2.4)	4 (0.9)	10 (2.7)	4 (1.3)	2 (1.8)
Total	441 (100)	412 (100)	427 (100)	374 (100)	306 (100)	111 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2015

Table 4. Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis (1), San Francisco, 2004-2015

Gender	Year of AIDS diagnosis					
	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)
Male	427 (87.9)	431 (89.4)	402 (89.7)	402 (90.1)	380 (86.8)	288 (88.6)
Female	41 (8.4)	37 (7.7)	31 (6.9)	32 (7.2)	44 (10.0)	25 (7.7)
Transgender	18 (3.7)	14 (2.9)	15 (3.3)	12 (2.7)	14 (3.2)	12 (3.7)
Total	486 (100)	482 (100)	448 (100)	446 (100)	438 (100)	325 (100)

Gender	Year of AIDS diagnosis					
	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)
Male	268 (89.6)	213 (84.9)	209 (87.1)	163 (88.1)	118 (90.1)	36 (81.8)
Female	16 (5.4)	24 (9.6)	25 (10.4)	17 (9.2)	6 (4.6)	7 (15.9)
Transgender	15 (5.0)	14 (5.6)	6 (2.5)	5 (2.7)	7 (5.3)	1 (2.3)
Total	299 (100)	251 (100)	240 (100)	185 (100)	131 (100)	44 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2015

Table 5. Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis (1), San Francisco, 2004-2015

Race/ethnicity	Year of initial HIV diagnosis					
	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)
White	392 (55.5)	294 (53.5)	279 (53.8)	269 (50.9)	256 (49.2)	238 (51.2)
African American	89 (12.6)	80 (14.5)	75 (14.5)	79 (14.9)	83 (16.0)	70 (15.1)
Latino	139 (19.7)	120 (21.8)	113 (21.8)	105 (19.8)	121 (23.3)	100 (21.5)
Asian/Pacific Islander/ Native American	58 (8.2)	36 (6.5)	36 (6.9)	51 (9.6)	44 (8.5)	37 (8.0)
Multiple races	26 (3.7)	18 (3.3)	16 (3.1)	23 (4.3)	16 (3.1)	20 (4.3)
Total (2)	706 (100)	550 (100)	519 (100)	529 (100)	520 (100)	465 (100)

Race/ethnicity	Year of initial HIV diagnosis					
	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)
White	212 (48.1)	213 (51.7)	213 (49.9)	171 (45.7)	135 (44.1)	46 (41.4)
African American	61 (13.8)	67 (16.3)	43 (10.1)	49 (13.1)	33 (10.8)	15 (13.5)
Latino	108 (24.5)	83 (20.1)	106 (24.8)	93 (24.9)	85 (27.8)	30 (27.0)
Asian/Pacific Islander/ Native American	38 (8.6)	33 (8.0)	54 (12.6)	51 (13.6)	40 (13.1)	16 (14.4)
Multiple races	21 (4.8)	14 (3.4)	9 (2.1)	8 (2.1)	13 (4.2)	4 (3.6)
Total (2)	441 (100)	412 (100)	427 (100)	374 (100)	306 (100)	111 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2015

Table 6. Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis (1), San Francisco, 2004-2015

Race/ethnicity	Year of AIDS diagnosis					
	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)
White	258 (53.1)	269 (55.8)	251 (56.0)	259 (58.1)	223 (50.9)	171 (52.6)
African American	76 (15.6)	82 (17.0)	78 (17.4)	75 (16.8)	81 (18.5)	63 (19.4)
Latino	115 (23.7)	95 (19.7)	79 (17.6)	75 (16.8)	76 (17.4)	65 (20.0)
Asian/Pacific Islander/ Native American	26 (5.3)	27 (5.6)	26 (5.8)	26 (5.8)	38 (8.7)	16 (4.9)
Multiple races	11 (2.3)	9 (1.9)	14 (3.1)	11 (2.5)	20 (4.6)	10 (3.1)
Total (2)	486 (100)	482 (100)	448 (100)	446 (100)	438 (100)	325 (100)

Race/ethnicity	Year of AIDS diagnosis					
	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)
White	147 (49.2)	131 (52.2)	122 (50.8)	90 (48.6)	48 (36.6)	19 (43.2)
African American	52 (17.4)	44 (17.5)	40 (16.7)	32 (17.3)	28 (21.4)	6 (13.6)
Latino	62 (20.7)	46 (18.3)	50 (20.8)	34 (18.4)	30 (22.9)	13 (29.5)
Asian/Pacific Islander/ Native American	26 (8.7)	15 (6.0)	17 (7.1)	19 (10.3)	16 (12.2)	3 (6.8)
Multiple races	12 (4.0)	14 (5.6)	11 (4.6)	10 (5.4)	9 (6.9)	3 (6.8)
Total (2)	299 (100)	251 (100)	240 (100)	185 (100)	131 (100)	44 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2015

Table 7. Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis (1), San Francisco, 2004-2015

Transmission category (2)	Year of initial HIV diagnosis					
	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)
Gay or bisexual male	484 (68.6)	376 (68.4)	365 (70.3)	346 (65.4)	373 (71.7)	326 (70.1)
Injection drug user (IDU)	63 (8.9)	36 (6.5)	39 (7.5)	34 (6.4)	33 (6.3)	25 (5.4)
Gay or bisexual male IDU	118 (16.7)	92 (16.7)	81 (15.6)	91 (17.2)	64 (12.3)	77 (16.6)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	21 (3.0)	30 (5.5)	24 (4.6)	42 (7.9)	36 (6.9)	23 (4.9)
Perinatal	2 (0.3)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	18 (2.5)	15 (2.7)	10 (1.9)	16 (3.0)	14 (2.7)	14 (3.0)
Total	706 (100)	550 (100)	519 (100)	529 (100)	520 (100)	465 (100)

Transmission category (2)	Year of initial HIV diagnosis					
	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)
Gay or bisexual male	284 (64.4)	297 (72.1)	332 (77.8)	287 (76.7)	231 (75.5)	82 (73.9)
Injection drug user (IDU)	35 (7.9)	28 (6.8)	14 (3.3)	21 (5.6)	20 (6.5)	7 (6.3)
Gay or bisexual male IDU	66 (15.0)	47 (11.4)	44 (10.3)	39 (10.4)	33 (10.8)	10 (9.0)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	35 (7.9)	26 (6.3)	24 (5.6)	19 (5.1)	8 (2.6)	5 (4.5)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	21 (4.8)	14 (3.4)	13 (3.0)	8 (2.1)	14 (4.6)	7 (6.3)
Total	441 (100)	412 (100)	427 (100)	374 (100)	306 (100)	111 (100)

- (1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.
- (2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2015

Table 8. Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis (1), San Francisco, 2004-2015

Transmission category (2)	Year of AIDS diagnosis					
	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)
Gay or bisexual male	315 (64.8)	295 (61.2)	287 (64.1)	286 (64.1)	262 (59.8)	209 (64.3)
Injection drug user (IDU)	51 (10.5)	51 (10.6)	40 (8.9)	37 (8.3)	35 (8.0)	36 (11.1)
Gay or bisexual male IDU	95 (19.5)	106 (22.0)	95 (21.2)	87 (19.5)	98 (22.4)	54 (16.6)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.3)
Heterosexual (3)	15 (3.1)	19 (3.9)	19 (4.2)	30 (6.7)	32 (7.3)	18 (5.5)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.3)
Other/Unknown	10 (2.1)	11 (2.3)	7 (1.6)	6 (1.3)	11 (2.5)	6 (1.8)
Total	486 (100)	482 (100)	448 (100)	446 (100)	438 (100)	325 (100)

Transmission category (2)	Year of AIDS diagnosis					
	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)
Gay or bisexual male	169 (56.5)	150 (59.8)	154 (64.2)	110 (59.5)	89 (67.9)	24 (54.5)
Injection drug user (IDU)	23 (7.7)	33 (13.1)	22 (9.2)	19 (10.3)	15 (11.5)	6 (13.6)
Gay or bisexual male IDU	83 (27.8)	47 (18.7)	37 (15.4)	43 (23.2)	19 (14.5)	7 (15.9)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	16 (5.4)	16 (6.4)	18 (7.5)	11 (5.9)	6 (4.6)	6 (13.6)
Perinatal	0 (0.0)	1 (0.4)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	8 (2.7)	4 (1.6)	9 (3.8)	2 (1.1)	2 (1.5)	1 (2.3)
Total	299 (100)	251 (100)	240 (100)	185 (100)	131 (100)	44 (100)

- (1) Data in recent years are incomplete due to delays in case reporting.
(2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2015

Table 9. Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2015

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with HIV infection (1)
2002 (2)	673	350	13403
2003	640	338	13705
2004	706	344	14067
2005	550	348	14269
2006	519	327	14461
2007	529	324	14666
2008	520	264	14922
2009	465	255	15132
2010	441	247	15326
2011	412	235	15503
2012	427	236	15694
2013	374	213	15855
2014	306	181	15980
2015	111	77	16014

(1) Data in recent years are incomplete due to delays in case/death reporting.

(2) Year of the mandatory reporting of HIV non-AIDS cases in California began.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2015

Table 10. Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2015

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with AIDS (1)
1980	3	0	3
1981	26	9	20
1982	99	32	87
1983	274	111	250
1984	558	273	535
1985	859	534	860
1986	1236	807	1289
1987	1632	878	2043
1988	1763	1040	2766
1989	2161	1279	3648
1990	2046	1364	4330
1991	2285	1512	5103
1992	2328	1641	5790
1993	2069	1603	6256
1994	1783	1601	6438
1995	1562	1485	6515
1996	1078	993	6600
1997	804	424	6980
1998	694	402	7272
1999	579	353	7498
2000	557	350	7705
2001	515	324	7896
2002	495	323	8068
2003	560	303	8325
2004	486	310	8501
2005	482	312	8671
2006	448	289	8830
2007	446	271	9005
2008	438	229	9214
2009	325	212	9327
2010	299	195	9431
2011	251	189	9493
2012	240	184	9549
2013	185	155	9579
2014	131	143	9567
2015	44	66	9545
Total	29741	20196	

(1) Data in recent years are incomplete due to delays in case/death reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2015

Table 11. Persons living with HIV infection by age, gender and race/ethnicity, San Francisco

Current age (years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander/ Native American No. (%)	Multiple races No. (%)	Total (1) No. (%)
Male						
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.3)	1 (0.0)
13 - 17	0 (0.0)	0 (0.0)	1 (0.0)	0 (0.0)	0 (0.0)	1 (0.0)
18 - 24	29 (0.3)	21 (1.4)	33 (1.3)	19 (2.1)	4 (1.3)	106 (0.7)
25 - 29	148 (1.6)	55 (3.6)	134 (5.1)	41 (4.6)	13 (4.1)	391 (2.6)
30 - 39	734 (7.8)	136 (8.8)	469 (17.9)	183 (20.6)	59 (18.7)	1583 (10.7)
40 - 49	2229 (23.8)	363 (23.6)	828 (31.5)	312 (35.1)	110 (34.8)	3847 (26.1)
50 - 59	3680 (39.3)	565 (36.7)	839 (31.9)	237 (26.6)	98 (31.0)	5423 (36.8)
60 +	2552 (27.2)	399 (25.9)	323 (12.3)	98 (11.0)	31 (9.8)	3403 (23.1)
Subtotal	9372 (100)	1539 (100)	2627 (100)	890 (100)	316 (100)	14755 (100)
Female						
0 - 12	0 (0.0)	1 (0.3)	1 (0.5)	0 (0.0)	0 (0.0)	2 (0.2)
13 - 17	0 (0.0)	0 (0.0)	1 (0.5)	0 (0.0)	0 (0.0)	1 (0.1)
18 - 24	2 (0.8)	4 (1.1)	8 (4.3)	1 (1.4)	3 (13.0)	18 (2.0)
25 - 29	6 (2.3)	8 (2.2)	7 (3.8)	2 (2.8)	0 (0.0)	23 (2.5)
30 - 39	34 (12.9)	34 (9.5)	34 (18.4)	15 (21.1)	7 (30.4)	125 (13.8)
40 - 49	78 (29.5)	82 (22.9)	42 (22.7)	22 (31.0)	6 (26.1)	230 (25.5)
50 - 59	100 (37.9)	151 (42.2)	54 (29.2)	22 (31.0)	7 (30.4)	335 (37.1)
60 +	44 (16.7)	78 (21.8)	38 (20.5)	9 (12.7)	0 (0.0)	169 (18.7)
Subtotal	264 (100)	358 (100)	185 (100)	71 (100)	23 (100)	903 (100)
Transgender	74 (100)	117 (100)	115 (100)	38 (100)	12 (100)	356 (100)
Total	9710 (100)	2014 (100)	2927 (100)	999 (100)	351 (100)	16014 (100)

(1) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2015

Table 12. Persons living with HIV infection by transmission category, gender and race/ethnicity, San Francisco

Transmission category (1)	White		African American		Latino		Asian/ Pacific Islander/ Native American		Multiple races		Total (2)	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Male												
Gay or bisexual male	7629	(81.4)	890	(57.8)	2144	(81.6)	750	(84.3)	216	(68.4)	11635	(78.9)
Injection drug user (IDU)	186	(2.0)	226	(14.7)	61	(2.3)	20	(2.2)	12	(3.8)	505	(3.4)
Gay or bisexual male IDU	1447	(15.4)	314	(20.4)	316	(12.0)	87	(9.8)	80	(25.3)	2244	(15.2)
Transfusion/Hemophiliac	6	(0.1)	3	(0.2)	2	(0.1)	3	(0.3)	0	(0.3)	14	(0.0)
Heterosexual (3)	32	(0.3)	68	(4.4)	53	(2.0)	13	(1.5)	3	(0.9)	169	(1.1)
Perinatal	1	(0.0)	2	(0.1)	2	(0.1)	2	(0.2)	2	(0.6)	9	(0.1)
Other/Unknown	71	(0.8)	36	(2.3)	49	(1.9)	15	(1.7)	3	(0.9)	179	(1.2)
Subtotal	9372	(100)	1539	(100)	2627	(100)	890	(100)	316	(100)	14755	(100)
Female												
Injection drug user (IDU)	157	(59.5)	192	(53.6)	59	(31.9)	16	(22.5)	14	(60.9)	438	(48.5)
Transfusion/Hemophilia	5	(1.9)	2	(0.6)	2	(1.1)	2	(2.8)	0	(0.0)	11	(1.2)
Heterosexual (3)	83	(31.4)	135	(37.7)	95	(51.4)	43	(60.6)	4	(17.4)	360	(39.9)
Perinatal	1	(0.4)	6	(1.7)	9	(4.9)	0	(0.0)	2	(8.7)	18	(2.0)
Other/Unknown	18	(6.8)	23	(6.4)	20	(10.8)	10	(14.1)	3	(13.0)	76	(8.4)
Subtotal	264	(100)	358	(100)	185	(100)	71	(100)	23	(100)	903	(100)
Transgender	74	(100)	117	(100)	115	(100)	38	(100)	12	(100)	356	(100)
Total	9710	(100)	2014	(100)	2927	(100)	999	(100)	351	(100)	16014	(100)

(1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.

(2) Total includes persons with unknown race.

(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.