

HIV SEMI-ANNUAL SURVEILLANCE REPORT

San Francisco Department of Public Health

HIV Cases Reported Through June 2018

Contents	Page
Surveillance Summary.....	1
Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	2
Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	2
Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis.....	3
Table 2: Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis.....	4
Table 3: Diagnoses of HIV infection by gender and year of initial HIV diagnosis	5
Table 4: Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis...	6
Table 5: Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis.....	7
Table 6: Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis.....	8
Table 7: Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis.....	9
Table 8: Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis.....	10
Table 9: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	11
Table 10: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	12
Table 11: Persons living with HIV infection by age, gender and race/ethnicity.....	13
Table 12: Persons living with HIV infection by transmission category, gender and race/ethnicity.....	14

The HIV Surveillance Report is published semi-annually by the
San Francisco Department of Public Health, Population Health Division
Applied Research, Community Health Epidemiology, and Surveillance Branch (ARCHES)
25 Van Ness Avenue, Suite 500, San Francisco, CA 94102; Phone (415) 437-6336, FAX (415) 431-0353

Director of Health: Barbara A. Garcia, MPA
Director of Population Health Division: Tomas Aragon, MD, DrPH
Director of ARCHES: Wayne Enanoria, PhD, MPH
Directors of HIV Surveillance Section: Susan Scheer, PhD, MPH; Ling Hsu, MPH

***The HIV Surveillance Report is accessible via internet:
<http://www.sfdph.org/dph/files/reports/default.asp>***

HIV SURVEILLANCE SUMMARY

San Francisco¹ (as of 06/30/2018)

Persons living with HIV infection ²	15,966
Diagnoses of HIV infection stage 3 (AIDS), cumulative	30,060
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	20,854

California³ (as of 12/31/2016)

Persons living with HIV infection ²	132,405
Diagnoses of HIV infection stage 3 (AIDS), cumulative	NA
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	NA

United States⁴ (as of 12/31/2015)

Persons living with HIV infection ²	991,289
Diagnoses of HIV infection stage 3 (AIDS), cumulative (as of 12/31/2016)	1,268,595
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	718,906


¹ Residents of San Francisco at time of HIV or AIDS diagnosis.

² Persons diagnosed with HIV infection regardless of disease stage.

³ CA data source: California HIV Surveillance Report 2016
https://www.cdph.ca.gov/Programs/CID/DOA/Pages/OA_case_surveillance_reports.aspx


⁴ US data source: CDC HIV Surveillance Report 2016, vol. 28 (Table 2b, 17b, 20b).
<https://www.cdc.gov/hiv/library/reports/hiv-surveillance.html>

Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2018*



* Reporting for recent years is incomplete. See Table 9 for actual numbers per year.

Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2018 †



† Reporting for recent years is incomplete. See Table 10 for actual numbers per year.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2018

Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis (1), San Francisco, 2007-2018

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	3 (0.6)	4 (0.8)	1 (0.2)	3 (0.7)	0 (0.0)	0 (0.0)
18 - 24	53 (9.9)	54 (10.4)	61 (13.0)	56 (12.1)	44 (10.4)	60 (13.0)
25 - 29	102 (19.1)	80 (15.4)	66 (14.0)	63 (13.7)	70 (16.5)	76 (16.5)
30 - 39	186 (34.8)	180 (34.7)	141 (29.9)	144 (31.2)	114 (26.8)	140 (30.4)
40 - 49	132 (24.7)	152 (29.3)	121 (25.7)	129 (28.0)	130 (30.6)	131 (28.4)
50 - 59	45 (8.4)	39 (7.5)	64 (13.6)	50 (10.8)	52 (12.2)	42 (9.1)
60 +	13 (2.4)	10 (1.9)	17 (3.6)	16 (3.5)	15 (3.5)	12 (2.6)
Total	534 (100)	519 (100)	471 (100)	461 (100)	425 (100)	461 (100)

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	1 (0.3)	3 (1.1)	0 (0.0)	3 (1.4)	0 (0.0)
18 - 24	57 (14.4)	37 (11.7)	37 (13.5)	31 (13.2)	23 (10.4)	13 (15.3)
25 - 29	79 (20.0)	57 (18.0)	63 (22.9)	56 (23.9)	40 (18.0)	19 (22.4)
30 - 39	113 (28.6)	91 (28.8)	84 (30.5)	75 (32.1)	74 (33.3)	23 (27.1)
40 - 49	98 (24.8)	75 (23.7)	57 (20.7)	37 (15.8)	45 (20.3)	19 (22.4)
50 - 59	35 (8.9)	41 (13.0)	18 (6.5)	24 (10.3)	26 (11.7)	6 (7.1)
60 +	13 (3.3)	14 (4.4)	13 (4.7)	11 (4.7)	11 (5.0)	5 (5.9)
Total	395 (100)	316 (100)	275 (100)	234 (100)	222 (100)	85 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2018

Table 2. Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis (1), San Francisco, 2007-2018

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.2)	1 (0.2)	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)
18 - 24	15 (3.4)	17 (3.9)	15 (4.6)	15 (5.0)	12 (4.8)	11 (4.5)
25 - 29	39 (8.8)	31 (7.1)	23 (7.1)	27 (9.0)	20 (8.0)	25 (10.3)
30 - 39	134 (30.1)	123 (28.3)	75 (23.1)	74 (24.6)	62 (24.7)	60 (24.7)
40 - 49	167 (37.5)	167 (38.4)	120 (37.0)	100 (33.2)	99 (39.4)	83 (34.2)
50 - 59	63 (14.2)	75 (17.2)	69 (21.3)	62 (20.6)	49 (19.5)	44 (18.1)
60 +	26 (5.8)	21 (4.8)	21 (6.5)	23 (7.6)	9 (3.6)	20 (8.2)
Total	445 (100)	435 (100)	324 (100)	301 (100)	251 (100)	243 (100)

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.8)	0 (0.0)
18 - 24	10 (5.3)	5 (3.6)	4 (3.2)	2 (2.0)	1 (0.8)	2 (4.7)
25 - 29	17 (8.9)	14 (10.0)	9 (7.2)	8 (8.1)	14 (11.1)	5 (11.6)
30 - 39	43 (22.6)	41 (29.3)	34 (27.2)	28 (28.3)	30 (23.8)	12 (27.9)
40 - 49	60 (31.6)	40 (28.6)	36 (28.8)	22 (22.2)	42 (33.3)	11 (25.6)
50 - 59	39 (20.5)	27 (19.3)	28 (22.4)	19 (19.2)	13 (10.3)	8 (18.6)
60 +	21 (11.1)	13 (9.3)	14 (11.2)	20 (20.2)	25 (19.8)	5 (11.6)
Total	190 (100)	140 (100)	125 (100)	99 (100)	126 (100)	43 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2018

Table 3. Diagnoses of HIV infection by gender and year of initial HIV diagnosis (1), San Francisco, 2007-2018

Gender	Year of initial HIV diagnosis					
	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)
Male	460 (86.1)	460 (88.6)	419 (89.0)	408 (88.5)	371 (87.3)	429 (93.1)
Female	49 (9.2)	40 (7.7)	30 (6.4)	36 (7.8)	41 (9.6)	23 (5.0)
Transgender	25 (4.7)	19 (3.7)	22 (4.7)	17 (3.7)	13 (3.1)	9 (2.0)
Total	534 (100)	519 (100)	471 (100)	461 (100)	425 (100)	461 (100)

Gender	Year of initial HIV diagnosis					
	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)
Male	361 (91.4)	293 (92.7)	242 (88.0)	203 (86.8)	190 (85.6)	71 (83.5)
Female	22 (5.6)	13 (4.1)	26 (9.5)	24 (10.3)	28 (12.6)	11 (12.9)
Transgender	12 (3.0)	10 (3.2)	7 (2.5)	7 (3.0)	4 (1.8)	3 (3.5)
Total	395 (100)	316 (100)	275 (100)	234 (100)	222 (100)	85 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2018

Table 4. Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis (1), San Francisco, 2007-2018

Gender	Year of AIDS diagnosis					
	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)
Male	401 (90.1)	375 (86.2)	288 (88.9)	269 (89.4)	213 (84.9)	210 (86.4)
Female	31 (7.0)	43 (9.9)	24 (7.4)	16 (5.3)	24 (9.6)	24 (9.9)
Transgender	13 (2.9)	17 (3.9)	12 (3.7)	16 (5.3)	14 (5.6)	9 (3.7)
Total	445 (100)	435 (100)	324 (100)	301 (100)	251 (100)	243 (100)

Gender	Year of AIDS diagnosis					
	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)
Male	167 (87.9)	126 (90.0)	100 (80.0)	86 (86.9)	100 (79.4)	37 (86.0)
Female	17 (8.9)	7 (5.0)	18 (14.4)	10 (10.1)	21 (16.7)	6 (14.0)
Transgender	6 (3.2)	7 (5.0)	7 (5.6)	3 (3.0)	5 (4.0)	0 (0.0)
Total	190 (100)	140 (100)	125 (100)	99 (100)	126 (100)	43 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2018

Table 5. Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis (1), San Francisco, 2007-2018

Race/ethnicity	Year of initial HIV diagnosis					
	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)
White	267 (50.0)	253 (48.7)	227 (48.2)	223 (48.4)	219 (51.5)	227 (49.2)
African American	81 (15.2)	80 (15.4)	63 (13.4)	62 (13.4)	65 (15.3)	47 (10.2)
Latino	107 (20.0)	118 (22.7)	113 (24.0)	107 (23.2)	85 (20.0)	114 (24.7)
Asian/Pacific Islander/ Native American	50 (9.4)	44 (8.5)	41 (8.7)	42 (9.1)	36 (8.5)	58 (12.6)
Multiple races	29 (5.4)	24 (4.6)	26 (5.5)	26 (5.6)	19 (4.5)	14 (3.0)
Total (2)	534 (100)	519 (100)	471 (100)	461 (100)	425 (100)	461 (100)

Race/ethnicity	Year of initial HIV diagnosis					
	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)
White	181 (45.8)	136 (43.0)	113 (41.1)	92 (39.3)	85 (38.3)	22 (25.9)
African American	48 (12.2)	32 (10.1)	40 (14.5)	34 (14.5)	38 (17.1)	21 (24.7)
Latino	100 (25.3)	88 (27.8)	73 (26.5)	67 (28.6)	56 (25.2)	31 (36.5)
Asian/Pacific Islander/ Native American	50 (12.7)	46 (14.6)	32 (11.6)	36 (15.4)	32 (14.4)	7 (8.2)
Multiple races	15 (3.8)	14 (4.4)	16 (5.8)	5 (2.1)	11 (5.0)	4 (4.7)
Total (2)	395 (100)	316 (100)	275 (100)	234 (100)	222 (100)	85 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2018

Table 6. Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis (1), San Francisco, 2007-2018

Race/ethnicity	Year of AIDS diagnosis					
	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)
White	252 (56.6)	219 (50.3)	171 (52.8)	145 (48.2)	128 (51.0)	121 (49.8)
African American	74 (16.6)	79 (18.2)	60 (18.5)	50 (16.6)	45 (17.9)	39 (16.0)
Latino	78 (17.5)	77 (17.7)	66 (20.4)	64 (21.3)	48 (19.1)	55 (22.6)
Asian/Pacific Islander/ Native American	26 (5.8)	39 (9.0)	15 (4.6)	25 (8.3)	14 (5.6)	15 (6.2)
Multiple races	15 (3.4)	21 (4.8)	12 (3.7)	17 (5.6)	16 (6.4)	13 (5.3)
Total (2)	445 (100)	435 (100)	324 (100)	301 (100)	251 (100)	243 (100)

Race/ethnicity	Year of AIDS diagnosis					
	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)
White	94 (49.5)	52 (37.1)	60 (48.0)	48 (48.5)	52 (41.3)	11 (25.6)
African American	32 (16.8)	29 (20.7)	19 (15.2)	15 (15.2)	23 (18.3)	9 (20.9)
Latino	34 (17.9)	31 (22.1)	31 (24.8)	20 (20.2)	34 (27.0)	15 (34.9)
Asian/Pacific Islander/ Native American	18 (9.5)	18 (12.9)	5 (4.0)	11 (11.1)	13 (10.3)	3 (7.0)
Multiple races	12 (6.3)	10 (7.1)	10 (8.0)	5 (5.1)	4 (3.2)	5 (11.6)
Total (2)	190 (100)	140 (100)	125 (100)	99 (100)	126 (100)	43 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2018

Table 7. Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis (1), San Francisco, 2007-2018

Transmission category (2)	Year of initial HIV diagnosis					
	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)
Gay or bisexual male	343 (64.2)	369 (71.1)	326 (69.2)	301 (65.3)	302 (71.1)	357 (77.4)
Injection drug user (IDU)	41 (7.7)	33 (6.4)	25 (5.3)	37 (8.0)	28 (6.6)	20 (4.3)
Gay or bisexual male IDU	91 (17.0)	70 (13.5)	81 (17.2)	68 (14.8)	59 (13.9)	48 (10.4)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	44 (8.2)	35 (6.7)	25 (5.3)	38 (8.2)	29 (6.8)	25 (5.4)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	15 (2.8)	12 (2.3)	14 (3.0)	17 (3.7)	7 (1.6)	11 (2.4)
Total	534 (100)	519 (100)	471 (100)	461 (100)	425 (100)	461 (100)

Transmission category (2)	Year of initial HIV diagnosis					
	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)
Gay or bisexual male	302 (76.5)	233 (73.7)	208 (75.6)	165 (70.5)	131 (59.0)	48 (56.5)
Injection drug user (IDU)	21 (5.3)	23 (7.3)	17 (6.2)	20 (8.5)	26 (11.7)	17 (20.0)
Gay or bisexual male IDU	48 (12.2)	42 (13.3)	24 (8.7)	23 (9.8)	32 (14.4)	7 (8.2)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	16 (4.1)	9 (2.8)	18 (6.5)	15 (6.4)	14 (6.3)	7 (8.2)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	8 (2.0)	9 (2.8)	8 (2.9)	11 (4.7)	19 (8.6)	6 (7.1)
Total	395 (100)	316 (100)	275 (100)	234 (100)	222 (100)	85 (100)

- (1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.
- (2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2018

Table 8. Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis (1), San Francisco, 2007-2018

Transmission category (2)	Year of AIDS diagnosis					
	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)
Gay or bisexual male	286 (64.3)	262 (60.2)	208 (64.2)	171 (56.8)	149 (59.4)	152 (62.6)
Injection drug user (IDU)	38 (8.5)	34 (7.8)	35 (10.8)	23 (7.6)	34 (13.5)	23 (9.5)
Gay or bisexual male IDU	89 (20.0)	98 (22.5)	56 (17.3)	84 (27.9)	47 (18.7)	42 (17.3)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	29 (6.5)	31 (7.1)	18 (5.6)	18 (6.0)	17 (6.8)	20 (8.2)
Perinatal	0 (0.0)	0 (0.0)	1 (0.3)	0 (0.0)	1 (0.4)	0 (0.0)
Other/Unknown	3 (0.7)	10 (2.3)	5 (1.5)	5 (1.7)	3 (1.2)	6 (2.5)
Total	445 (100)	435 (100)	324 (100)	301 (100)	251 (100)	243 (100)

Transmission category (2)	Year of AIDS diagnosis					
	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)
Gay or bisexual male	112 (58.9)	95 (67.9)	71 (56.8)	58 (58.6)	59 (46.8)	22 (51.2)
Injection drug user (IDU)	19 (10.0)	16 (11.4)	18 (14.4)	13 (13.1)	13 (10.3)	6 (14.0)
Gay or bisexual male IDU	46 (24.2)	21 (15.0)	23 (18.4)	17 (17.2)	29 (23.0)	9 (20.9)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	12 (6.3)	6 (4.3)	10 (8.0)	9 (9.1)	19 (15.1)	4 (9.3)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	1 (0.5)	2 (1.4)	3 (2.4)	2 (2.0)	6 (4.8)	2 (4.7)
Total	190 (100)	140 (100)	125 (100)	99 (100)	126 (100)	43 (100)

- (1) Data in recent years are incomplete due to delays in case reporting.
(2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2018

Table 9. Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2018

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with HIV infection (1)
2002 (2)	664	349	13347
2003	655	338	13664
2004	688	342	14010
2005	568	348	14230
2006	533	327	14436
2007	534	325	14645
2008	519	265	14899
2009	471	254	15116
2010	461	246	15331
2011	425	235	15521
2012	461	237	15745
2013	395	261	15879
2014	316	241	15954
2015	275	256	15973
2016	234	235	15972
2017	222	244	15950
2018	85	69	15966

(1) Data in recent years are incomplete due to delays in case/death reporting.

(2) Year of the mandatory reporting of HIV non-AIDS cases in California began.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2018

Table 10. Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2018

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with AIDS (1)
1980	3	0	3
1981	26	8	21
1982	99	32	88
1983	274	111	251
1984	557	273	535
1985	860	534	861
1986	1236	807	1290
1987	1632	878	2044
1988	1763	1040	2767
1989	2161	1279	3649
1990	2045	1365	4329
1991	2285	1512	5102
1992	2328	1640	5790
1993	2067	1603	6254
1994	1779	1600	6433
1995	1555	1485	6503
1996	1076	995	6584
1997	800	424	6960
1998	693	402	7251
1999	577	353	7475
2000	556	350	7681
2001	511	324	7868
2002	492	323	8037
2003	553	303	8287
2004	480	308	8459
2005	479	312	8626
2006	451	289	8788
2007	445	271	8962
2008	435	229	9168
2009	324	211	9281
2010	301	194	9388
2011	251	189	9450
2012	243	183	9510
2013	190	192	9508
2014	140	189	9459
2015	125	203	9381
2016	99	186	9294
2017	126	196	9224
2018	43	61	9206
Total	30060	20854	

(1) Data in recent years are incomplete due to delays in case/death reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2018

Table 11. Persons living with HIV infection by age, gender and race/ethnicity, San Francisco

Current age (years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander/ Native American No. (%)	Multiple races No. (%)	Total (1) No. (%)
Male						
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.2)	1 (0.0)
18 - 24	20 (0.2)	18 (1.2)	23 (0.8)	7 (0.7)	3 (0.7)	71 (0.5)
25 - 29	100 (1.1)	49 (3.4)	121 (4.3)	50 (5.3)	13 (2.9)	333 (2.3)
30 - 39	641 (7.1)	146 (10.1)	466 (16.7)	194 (20.6)	67 (14.7)	1514 (10.3)
40 - 49	1553 (17.2)	255 (17.6)	727 (26.1)	286 (30.3)	126 (27.7)	2950 (20.1)
50 - 59	3523 (39.0)	495 (34.1)	978 (35.1)	275 (29.2)	150 (33.0)	5422 (37.0)
60 +	3186 (35.3)	488 (33.6)	474 (17.0)	131 (13.9)	95 (20.9)	4376 (29.8)
Subtotal	9023 (100)	1451 (100)	2789 (100)	943 (100)	455 (100)	14667 (100)
Female						
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	1 (0.3)	1 (0.5)	0 (0.0)	0 (0.0)	2 (0.2)
18 - 24	3 (1.1)	8 (2.4)	5 (2.5)	0 (0.0)	4 (11.4)	20 (2.2)
25 - 29	6 (2.3)	6 (1.8)	9 (4.5)	3 (4.2)	0 (0.0)	24 (2.6)
30 - 39	28 (10.7)	35 (10.3)	39 (19.6)	10 (14.1)	4 (11.4)	116 (12.8)
40 - 49	63 (24.0)	56 (16.5)	40 (20.1)	23 (32.4)	11 (31.4)	194 (21.3)
50 - 59	102 (38.9)	126 (37.1)	54 (27.1)	21 (29.6)	12 (34.3)	315 (34.7)
60 +	60 (22.9)	108 (31.8)	51 (25.6)	14 (19.7)	4 (11.4)	238 (26.2)
Subtotal	262 (100)	340 (100)	199 (100)	71 (100)	35 (100)	909 (100)
Transgender	72 (100)	124 (100)	135 (100)	40 (100)	19 (100)	390 (100)
Total	9357 (100)	1915 (100)	3123 (100)	1054 (100)	509 (100)	15966 (100)

(1) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2018

Table 12. Persons living with HIV infection by transmission category, gender and race/ethnicity, San Francisco

Transmission category (1)	White		African American		Latino		Asian/ Pacific Islander/ Native American		Multiple races		Total (2)	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Male												
Gay or bisexual male	7363	(81.6)	860	(59.3)	2289	(82.1)	804	(85.3)	328	(72.1)	11645	(79.4)
Injection drug user (IDU)	173	(1.9)	192	(13.2)	69	(2.5)	20	(2.1)	17	(3.7)	471	(3.2)
Gay or bisexual male IDU	1397	(15.5)	292	(20.1)	329	(11.8)	79	(8.4)	98	(21.5)	2195	(15.0)
Transfusion/Hemophiliac	6	(0.1)	2	(0.1)	2	(0.1)	3	(0.3)	1	(0.3)	14	(0.2)
Heterosexual (3)	29	(0.3)	66	(4.5)	53	(1.9)	20	(2.1)	6	(1.3)	174	(1.2)
Perinatal	1	(0.0)	3	(0.2)	2	(0.1)	1	(0.1)	2	(0.4)	9	(0.1)
Other/Unknown	54	(0.6)	36	(2.5)	45	(1.6)	16	(1.7)	3	(0.7)	159	(1.1)
Subtotal	9023	(100)	1451	(100)	2789	(100)	943	(100)	455	(100)	14667	(100)
Female												
Injection drug user (IDU)	147	(56.1)	170	(50.0)	60	(30.2)	13	(18.3)	24	(68.6)	414	(45.5)
Transfusion/Hemophilia	5	(1.9)	2	(0.6)	2	(1.0)	2	(2.8)	0	(0.0)	11	(1.2)
Heterosexual (3)	87	(33.2)	138	(40.6)	108	(54.3)	50	(70.4)	7	(20.0)	390	(42.9)
Perinatal	1	(0.4)	6	(1.8)	9	(4.5)	0	(0.0)	2	(5.7)	18	(2.0)
Other/Unknown	22	(8.4)	24	(7.1)	20	(10.1)	6	(8.5)	2	(5.7)	76	(8.4)
Subtotal	262	(100)	340	(100)	199	(100)	71	(100)	35	(100)	909	(100)
Transgender	72	(100)	124	(100)	135	(100)	40	(100)	19	(100)	390	(100)
Total	9357	(100)	1915	(100)	3123	(100)	1054	(100)	509	(100)	15966	(100)

(1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.

(2) Total includes persons with unknown race.

(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.