

HIV SEMI-ANNUAL SURVEILLANCE REPORT

San Francisco Department of Public Health

HIV Cases Reported Through June 2016

Contents	Page
Surveillance Summary.....	1
Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	2
Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	2
Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis.....	3
Table 2: Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis.....	4
Table 3: Diagnoses of HIV infection by gender and year of initial HIV diagnosis	5
Table 4: Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis...	6
Table 5: Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis.....	7
Table 6: Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis.....	8
Table 7: Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis.....	9
Table 8: Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis.....	10
Table 9: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	11
Table 10: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	12
Table 11: Persons living with HIV infection by age, gender and race/ethnicity.....	13
Table 12: Persons living with HIV infection by transmission category, gender and race/ethnicity.....	14

The HIV Surveillance Report is published semi-annually by the
San Francisco Department of Public Health, Population Health Division
Applied Research, Community Health Epidemiology, and Surveillance Branch (ARCHES)
25 Van Ness Avenue, Suite 500, San Francisco, CA 94102; Phone (415) 437-6336, FAX (415) 431-0353

Director of Health: Barbara A. Garcia, MPA
Director of Population Health Division: Tomas Aragon, MD, DrPH
HIV Surveillance Section Directors: Susan Scheer, PhD, MPH; Ling Hsu, MPH

***The HIV Surveillance Report is accessible via internet:
<http://www.sfdph.org/dph/files/reports/default.asp>***

HIV SURVEILLANCE SUMMARY

San Francisco¹ (as of 06/30/2016)

Persons living with HIV infection ²	16,030
Diagnoses of HIV infection stage 3 (AIDS), cumulative	29,850
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	20,422

California³ (as of 12/31/2013)

Persons living with HIV infection ²	119,878
Diagnoses of HIV infection stage 3 (AIDS), cumulative	169,734
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	96,443

United States⁴ (as of 12/31/2013)

Persons living with HIV infection ²	949,331
Diagnoses of HIV infection stage 3 (AIDS), cumulative (as of 12/31/2014)	1,236,994
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	691,458


¹ Residents of San Francisco at time of HIV or AIDS diagnosis.

² Persons diagnosed with HIV infection regardless of disease stage.

³ CA data source: www.cdph.ca.gov/data/statistics/Pages/OAHIVAIDSStatistics.aspx.


⁴ US data source: CDC HIV Surveillance Report 2014, vol. 26 (Table 2b, 13b, 18b).
<http://www.cdc.gov/hiv/library/reports/surveillance/>.

Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2016*



* Reporting for recent years is incomplete. See Table 9 for actual numbers per year.

Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2016 †



† Reporting for recent years is incomplete. See Table 10 for actual numbers per year.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2016

Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis (1), San Francisco, 2005-2016

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)
0 - 12	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.2)	0 (0.0)	2 (0.4)	4 (0.8)	1 (0.2)	3 (0.7)
18 - 24	56 (9.8)	65 (12.3)	54 (10.2)	55 (10.7)	57 (12.2)	53 (11.7)
25 - 29	76 (13.3)	61 (11.5)	101 (19.0)	78 (15.1)	65 (13.9)	63 (13.9)
30 - 39	214 (37.5)	181 (34.2)	186 (35.0)	182 (35.3)	143 (30.5)	142 (31.4)
40 - 49	160 (28.0)	149 (28.2)	130 (24.5)	150 (29.1)	123 (26.2)	126 (27.9)
50 - 59	50 (8.8)	55 (10.4)	45 (8.5)	40 (7.8)	63 (13.4)	49 (10.8)
60 +	13 (2.3)	18 (3.4)	13 (2.4)	7 (1.4)	17 (3.6)	16 (3.5)
Total	571 (100)	529 (100)	531 (100)	516 (100)	469 (100)	452 (100)

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.2)	0 (0.0)	0 (0.0)	1 (0.3)	3 (1.1)	0 (0.0)
18 - 24	44 (10.5)	58 (12.8)	54 (14.0)	36 (11.7)	35 (13.4)	14 (14.1)
25 - 29	65 (15.5)	72 (15.9)	76 (19.7)	53 (17.2)	59 (22.5)	22 (22.2)
30 - 39	111 (26.5)	139 (30.6)	111 (28.8)	92 (29.9)	78 (29.8)	29 (29.3)
40 - 49	132 (31.5)	130 (28.6)	96 (24.9)	74 (24.0)	58 (22.1)	19 (19.2)
50 - 59	52 (12.4)	43 (9.5)	35 (9.1)	39 (12.7)	17 (6.5)	12 (12.1)
60 +	14 (3.3)	12 (2.6)	13 (3.4)	13 (4.2)	12 (4.6)	3 (3.0)
Total	419 (100)	454 (100)	385 (100)	308 (100)	262 (100)	99 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report

Cases Reported as of 06/30/2016

Table 2. Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis (1), San Francisco, 2005-2016

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	0 (0.0)	1 (0.2)	1 (0.2)	1 (0.3)	0 (0.0)
18 - 24	14 (2.9)	13 (2.9)	15 (3.4)	17 (3.9)	15 (4.6)	15 (5.0)
25 - 29	32 (6.7)	25 (5.6)	39 (8.7)	31 (7.1)	23 (7.1)	27 (9.0)
30 - 39	157 (32.7)	125 (27.8)	133 (29.8)	124 (28.5)	75 (23.1)	73 (24.4)
40 - 49	183 (38.1)	184 (40.9)	167 (37.4)	166 (38.2)	121 (37.2)	99 (33.1)
50 - 59	80 (16.7)	82 (18.2)	65 (14.6)	75 (17.2)	69 (21.2)	62 (20.7)
60 +	14 (2.9)	21 (4.7)	26 (5.8)	21 (4.8)	21 (6.5)	23 (7.7)
Total	480 (100)	450 (100)	446 (100)	435 (100)	325 (100)	299 (100)

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
18 - 24	12 (4.8)	11 (4.6)	10 (5.3)	4 (2.9)	4 (3.3)	1 (2.7)
25 - 29	20 (8.0)	25 (10.4)	17 (9.0)	14 (10.2)	8 (6.6)	2 (5.4)
30 - 39	62 (24.8)	60 (24.9)	43 (22.8)	40 (29.2)	33 (27.0)	8 (21.6)
40 - 49	98 (39.2)	80 (33.2)	59 (31.2)	39 (28.5)	36 (29.5)	11 (29.7)
50 - 59	49 (19.6)	45 (18.7)	39 (20.6)	26 (19.0)	27 (22.1)	5 (13.5)
60 +	9 (3.6)	20 (8.3)	21 (11.1)	14 (10.2)	14 (11.5)	10 (27.0)
Total	250 (100)	241 (100)	189 (100)	137 (100)	122 (100)	37 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2016

Table 3. Diagnoses of HIV infection by gender and year of initial HIV diagnosis (1), San Francisco, 2005-2016

Gender	Year of initial HIV diagnosis					
	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)
Male	510 (89.3)	480 (90.7)	459 (86.4)	460 (89.1)	415 (88.5)	402 (88.9)
Female	45 (7.9)	36 (6.8)	48 (9.0)	39 (7.6)	30 (6.4)	35 (7.7)
Transgender	16 (2.8)	13 (2.5)	24 (4.5)	17 (3.3)	24 (5.1)	15 (3.3)
Total	571 (100)	529 (100)	531 (100)	516 (100)	469 (100)	452 (100)

Gender	Year of initial HIV diagnosis					
	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)
Male	363 (86.6)	425 (93.6)	353 (91.7)	290 (94.2)	231 (88.2)	88 (88.9)
Female	41 (9.8)	23 (5.1)	22 (5.7)	12 (3.9)	24 (9.2)	10 (10.1)
Transgender	15 (3.6)	6 (1.3)	10 (2.6)	6 (1.9)	7 (2.7)	1 (1.0)
Total	419 (100)	454 (100)	385 (100)	308 (100)	262 (100)	99 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2016

Table 4. Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis (1), San Francisco, 2005-2016

Gender	Year of AIDS diagnosis					
	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)
Male	428 (89.2)	402 (89.3)	402 (90.1)	376 (86.4)	289 (88.9)	267 (89.3)
Female	38 (7.9)	32 (7.1)	32 (7.2)	43 (9.9)	24 (7.4)	16 (5.4)
Transgender	14 (2.9)	16 (3.6)	12 (2.7)	16 (3.7)	12 (3.7)	16 (5.4)
Total	480 (100)	450 (100)	446 (100)	435 (100)	325 (100)	299 (100)

Gender	Year of AIDS diagnosis					
	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)
Male	212 (84.8)	209 (86.7)	167 (88.4)	122 (89.1)	100 (82.0)	34 (91.9)
Female	24 (9.6)	24 (10.0)	17 (9.0)	7 (5.1)	17 (13.9)	2 (5.4)
Transgender	14 (5.6)	8 (3.3)	5 (2.6)	8 (5.8)	5 (4.1)	1 (2.7)
Total	250 (100)	241 (100)	189 (100)	137 (100)	122 (100)	37 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2016

Table 5. Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis (1), San Francisco, 2005-2016

Race/ethnicity	Year of initial HIV diagnosis					
	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)
White	303 (53.1)	288 (54.4)	272 (51.2)	255 (49.4)	228 (48.6)	219 (48.5)
African American	80 (14.0)	76 (14.4)	79 (14.9)	80 (15.5)	66 (14.1)	63 (13.9)
Latino	127 (22.2)	113 (21.4)	105 (19.8)	119 (23.1)	112 (23.9)	108 (23.9)
Asian/Pacific Islander/ Native American	37 (6.5)	35 (6.6)	49 (9.2)	42 (8.1)	41 (8.7)	40 (8.8)
Multiple races	23 (4.0)	17 (3.2)	26 (4.9)	20 (3.9)	21 (4.5)	21 (4.6)
Total (2)	571 (100)	529 (100)	531 (100)	516 (100)	469 (100)	452 (100)

Race/ethnicity	Year of initial HIV diagnosis					
	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)
White	221 (52.7)	227 (50.0)	174 (45.2)	133 (43.2)	110 (42.0)	42 (42.4)
African American	65 (15.5)	46 (10.1)	49 (12.7)	32 (10.4)	44 (16.8)	13 (13.1)
Latino	82 (19.6)	110 (24.2)	98 (25.5)	85 (27.6)	68 (26.0)	28 (28.3)
Asian/Pacific Islander/ Native American	35 (8.4)	58 (12.8)	50 (13.0)	43 (14.0)	32 (12.2)	13 (13.1)
Multiple races	15 (3.6)	12 (2.6)	12 (3.1)	15 (4.9)	7 (2.7)	2 (2.0)
Total (2)	419 (100)	454 (100)	385 (100)	308 (100)	262 (100)	99 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2016

Table 6. Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis (1), San Francisco, 2005-2016

Race/ethnicity	Year of AIDS diagnosis					
	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)
White	267 (55.6)	249 (55.3)	257 (57.6)	220 (50.6)	171 (52.6)	145 (48.5)
African American	82 (17.1)	81 (18.0)	75 (16.8)	80 (18.4)	62 (19.1)	52 (17.4)
Latino	94 (19.6)	79 (17.6)	75 (16.8)	76 (17.5)	66 (20.3)	62 (20.7)
Asian/Pacific Islander/ Native American	26 (5.4)	26 (5.8)	26 (5.8)	39 (9.0)	16 (4.9)	25 (8.4)
Multiple races	11 (2.3)	15 (3.3)	13 (2.9)	20 (4.6)	10 (3.1)	15 (5.0)
Total (2)	480 (100)	450 (100)	446 (100)	435 (100)	325 (100)	299 (100)

Race/ethnicity	Year of AIDS diagnosis					
	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)
White	131 (52.4)	122 (50.6)	93 (49.2)	51 (37.2)	57 (46.7)	17 (45.9)
African American	44 (17.6)	38 (15.8)	32 (16.9)	29 (21.2)	19 (15.6)	7 (18.9)
Latino	46 (18.4)	52 (21.6)	34 (18.0)	30 (21.9)	32 (26.2)	9 (24.3)
Asian/Pacific Islander/ Native American	15 (6.0)	17 (7.1)	19 (10.1)	17 (12.4)	5 (4.1)	3 (8.1)
Multiple races	14 (5.6)	12 (5.0)	11 (5.8)	10 (7.3)	8 (6.6)	1 (2.7)
Total (2)	250 (100)	241 (100)	189 (100)	137 (100)	122 (100)	37 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2016

Table 7. Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis (1), San Francisco, 2005-2016

Transmission category (2)	Year of initial HIV diagnosis					
	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)
Gay or bisexual male	398 (69.7)	365 (69.0)	344 (64.8)	371 (71.9)	324 (69.1)	291 (64.4)
Injection drug user (IDU)	36 (6.3)	37 (7.0)	40 (7.5)	34 (6.6)	25 (5.3)	36 (8.0)
Gay or bisexual male IDU	91 (15.9)	91 (17.2)	90 (16.9)	65 (12.6)	78 (16.6)	69 (15.3)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	30 (5.3)	27 (5.1)	43 (8.1)	34 (6.6)	26 (5.5)	36 (8.0)
Perinatal	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	15 (2.6)	9 (1.7)	14 (2.6)	12 (2.3)	16 (3.4)	20 (4.4)
Total	571 (100)	529 (100)	531 (100)	516 (100)	469 (100)	452 (100)

Transmission category (2)	Year of initial HIV diagnosis					
	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)
Gay or bisexual male	300 (71.6)	353 (77.8)	299 (77.7)	229 (74.4)	188 (71.8)	71 (71.7)
Injection drug user (IDU)	28 (6.7)	18 (4.0)	21 (5.5)	22 (7.1)	17 (6.5)	9 (9.1)
Gay or bisexual male IDU	53 (12.6)	45 (9.9)	42 (10.9)	37 (12.0)	26 (9.9)	8 (8.1)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	27 (6.4)	25 (5.5)	16 (4.2)	8 (2.6)	16 (6.1)	7 (7.1)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	11 (2.6)	13 (2.9)	7 (1.8)	12 (3.9)	15 (5.7)	4 (4.0)
Total	419 (100)	454 (100)	385 (100)	308 (100)	262 (100)	99 (100)

- (1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.
- (2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report

Cases Reported as of 06/30/2016

Table 8. Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis (1), San Francisco, 2005-2016

Transmission category (2)	Year of AIDS diagnosis					
	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)
Gay or bisexual male	294 (61.3)	286 (63.6)	288 (64.6)	262 (60.2)	210 (64.6)	169 (56.5)
Injection drug user (IDU)	50 (10.4)	39 (8.7)	38 (8.5)	34 (7.8)	35 (10.8)	23 (7.7)
Gay or bisexual male IDU	106 (22.1)	98 (21.8)	87 (19.5)	97 (22.3)	54 (16.6)	83 (27.8)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.3)	0 (0.0)
Heterosexual (3)	19 (4.0)	21 (4.7)	29 (6.5)	32 (7.4)	18 (5.5)	17 (5.7)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.3)	0 (0.0)
Other/Unknown	11 (2.3)	6 (1.3)	4 (0.9)	10 (2.3)	6 (1.8)	7 (2.3)
Total	480 (100)	450 (100)	446 (100)	435 (100)	325 (100)	299 (100)

Transmission category (2)	Year of AIDS diagnosis					
	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)
Gay or bisexual male	149 (59.6)	155 (64.3)	112 (59.3)	93 (67.9)	68 (55.7)	25 (67.6)
Injection drug user (IDU)	33 (13.2)	21 (8.7)	19 (10.1)	15 (10.9)	18 (14.8)	4 (10.8)
Gay or bisexual male IDU	47 (18.8)	38 (15.8)	45 (23.8)	20 (14.6)	22 (18.0)	4 (10.8)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	17 (6.8)	20 (8.3)	11 (5.8)	6 (4.4)	9 (7.4)	4 (10.8)
Perinatal	1 (0.4)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	3 (1.2)	7 (2.9)	2 (1.1)	3 (2.2)	5 (4.1)	0 (0.0)
Total	250 (100)	241 (100)	189 (100)	137 (100)	122 (100)	37 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.

(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2016

Table 9. Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2016

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with HIV infection (1)
2002 (2)	664	350	13364
2003	655	338	13681
2004	688	343	14026
2005	571	348	14249
2006	529	328	14450
2007	531	324	14657
2008	516	264	14909
2009	469	254	15124
2010	452	247	15329
2011	419	234	15514
2012	454	237	15731
2013	385	261	15855
2014	308	217	15946
2015	262	199	16009
2016	99	78	16030

(1) Data in recent years are incomplete due to delays in case/death reporting.

(2) Year of the mandatory reporting of HIV non-AIDS cases in California began.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2016

Table 10. Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2016

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with AIDS (1)
1980	3	0	3
1981	26	8	21
1982	99	32	88
1983	274	111	251
1984	558	273	536
1985	860	534	862
1986	1236	807	1291
1987	1632	878	2045
1988	1763	1040	2768
1989	2162	1279	3651
1990	2046	1365	4332
1991	2286	1512	5106
1992	2329	1641	5794
1993	2069	1603	6260
1994	1784	1601	6443
1995	1557	1485	6515
1996	1076	993	6598
1997	803	424	6977
1998	694	402	7269
1999	577	353	7493
2000	556	350	7699
2001	514	324	7889
2002	493	323	8059
2003	558	303	8314
2004	484	309	8489
2005	480	312	8657
2006	450	289	8818
2007	446	271	8993
2008	435	229	9199
2009	325	211	9313
2010	299	195	9417
2011	250	188	9479
2012	241	184	9536
2013	189	192	9533
2014	137	171	9499
2015	122	162	9459
2016	37	68	9428
Total	29850	20422	

(1) Data in recent years are incomplete due to delays in case/death reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2016

Table 11. Persons living with HIV infection by age, gender and race/ethnicity, San Francisco

Current age (years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander/ Native American No. (%)	Multiple races No. (%)	Total (1) No. (%)
Male						
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.3)	1 (0.0)
13 - 17	0 (0.0)	1 (0.1)	1 (0.0)	0 (0.0)	0 (0.0)	2 (0.0)
18 - 24	24 (0.3)	26 (1.7)	25 (0.9)	21 (2.3)	4 (1.1)	100 (0.7)
25 - 29	126 (1.4)	51 (3.4)	139 (5.2)	43 (4.7)	12 (3.3)	371 (2.5)
30 - 39	708 (7.6)	139 (9.2)	485 (18.0)	179 (19.7)	63 (17.5)	1576 (10.7)
40 - 49	1963 (21.2)	315 (20.9)	785 (29.1)	305 (33.5)	110 (30.5)	3481 (23.6)
50 - 59	3676 (39.7)	537 (35.6)	897 (33.3)	253 (27.8)	118 (32.7)	5483 (37.2)
60 +	2772 (29.9)	439 (29.1)	362 (13.4)	109 (12.0)	53 (14.7)	3736 (25.3)
Subtotal	9269 (100)	1508 (100)	2694 (100)	910 (100)	361 (100)	14750 (100)
Female						
0 - 12	0 (0.0)	1 (0.3)	1 (0.5)	0 (0.0)	0 (0.0)	2 (0.2)
13 - 17	0 (0.0)	0 (0.0)	1 (0.5)	0 (0.0)	0 (0.0)	1 (0.1)
18 - 24	2 (0.8)	5 (1.4)	7 (3.7)	1 (1.4)	3 (11.1)	18 (2.0)
25 - 29	8 (3.0)	8 (2.3)	8 (4.2)	2 (2.9)	0 (0.0)	26 (2.9)
30 - 39	34 (12.8)	31 (8.9)	37 (19.4)	13 (18.8)	6 (22.2)	121 (13.4)
40 - 49	65 (24.4)	75 (21.5)	42 (22.0)	21 (30.4)	10 (37.0)	214 (23.7)
50 - 59	109 (41.0)	144 (41.3)	52 (27.2)	22 (31.9)	7 (25.9)	335 (37.1)
60 +	48 (18.0)	85 (24.4)	43 (22.5)	10 (14.5)	1 (3.7)	187 (20.7)
Subtotal	266 (100)	349 (100)	191 (100)	69 (100)	27 (100)	904 (100)
Transgender	75 (100)	125 (100)	122 (100)	39 (100)	15 (100)	376 (100)
Total	9610 (100)	1982 (100)	3007 (100)	1018 (100)	403 (100)	16030 (100)

(1) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2016

Table 12. Persons living with HIV infection by transmission category, gender and race/ethnicity, San Francisco

Transmission category (1)	White		African American		Latino		Asian/ Pacific Islander/ Native American		Multiple races		Total (2)	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Male												
Gay or bisexual male	7550	(81.5)	886	(58.8)	2202	(81.7)	773	(84.9)	250	(69.3)	11664	(79.1)
Injection drug user (IDU)	186	(2.0)	210	(13.9)	65	(2.4)	20	(2.2)	14	(3.9)	495	(3.4)
Gay or bisexual male IDU	1430	(15.4)	301	(20.0)	323	(12.0)	81	(8.9)	87	(24.1)	2222	(15.1)
Transfusion/Hemophiliac	6	(0.1)	2	(0.1)	2	(0.1)	3	(0.3)	1	(0.3)	14	(0.3)
Heterosexual (3)	30	(0.3)	69	(4.6)	53	(2.0)	14	(1.5)	4	(1.1)	170	(1.2)
Perinatal	1	(0.0)	3	(0.2)	2	(0.1)	1	(0.1)	2	(0.6)	9	(0.1)
Other/Unknown	66	(0.7)	37	(2.5)	47	(1.7)	18	(2.0)	3	(0.8)	176	(1.2)
Subtotal	9269	(100)	1508	(100)	2694	(100)	910	(100)	361	(100)	14750	(100)
Female												
Injection drug user (IDU)	150	(56.4)	185	(53.0)	59	(30.9)	15	(21.7)	17	(63.0)	426	(47.1)
Transfusion/Hemophilia	5	(1.9)	2	(0.6)	2	(1.0)	2	(2.9)	0	(0.0)	11	(1.2)
Heterosexual (3)	90	(33.8)	136	(39.0)	101	(52.9)	44	(63.8)	6	(22.2)	377	(41.7)
Perinatal	1	(0.4)	6	(1.7)	9	(4.7)	0	(0.0)	2	(7.4)	18	(2.0)
Other/Unknown	20	(7.5)	20	(5.7)	20	(10.5)	8	(11.6)	2	(7.4)	72	(8.0)
Subtotal	266	(100)	349	(100)	191	(100)	69	(100)	27	(100)	904	(100)
Transgender	75	(100)	125	(100)	122	(100)	39	(100)	15	(100)	376	(100)
Total	9610	(100)	1982	(100)	3007	(100)	1018	(100)	403	(100)	16030	(100)

- (1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (2) Total includes persons with unknown race.
- (3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.