

QUARTERLY AIDS SURVEILLANCE REPORT

San Francisco Department of Public Health AIDS Cases Reported Through June 2007

Contents	Page
Surveillance Summary.....	1
Table 1: Adult/Adolescent AIDS Cases by Transmission Category.....	2
Table 2: AIDS Cases by Gender and Year of Diagnosis.....	2
Table 3: AIDS Cases by Transmission Category and Race/Ethnicity.....	3
Table 4: AIDS Cases by Transmission Category and Asian/Pacific Islander Ethnicity.....	3
Table 5: AIDS Cases by Transmission Category and Year of Diagnosis.....	4
Table 6: AIDS Cases by Gender, Age Group and Race/Ethnicity.....	5
Table 7: AIDS Cases by Race/Ethnicity and Year of Diagnosis.....	6
Table 8: AIDS Cases and Cumulative Rates per 100,000 by Race/Ethnicity and Gender.....	6
Table 9: AIDS Incidence, Mortality, and Prevalence by Year.....	7
Table 10: Cases by Initial AIDS-Defining Condition.....	8
Table 11: Cumulative AIDS Indicator Conditions among Persons with AIDS.....	9
Table 12: Living Adult/Adolescent AIDS Cases by Transmission Category.....	10
Table 13: Living AIDS Cases by Transmission Category and Race/Ethnicity.....	10
Table 14: Living AIDS Cases by Gender, Age Group and Race/Ethnicity.....	11
Table 15: Living AIDS Cases by Initial AIDS-Defining Condition.....	12
Table 16: Cumulative AIDS Indicator Conditions among Persons Living with AIDS.....	13

The AIDS Surveillance Report is published quarterly by the
San Francisco Department of Public Health, HIV Epidemiology Section
25 Van Ness Avenue, Suite 500, San Francisco, CA 94102; Phone (415) 554-9050, FAX (415) 431-0353
Director of Health: Mitchell Katz, MD; Section Co-Directors: Ling Hsu, MPH, Susan Scheer, PhD, MPH;
Program Coordinators: Maree Kay Parisi, Viva Delgado, MPH;
Epidemiologists: Anne Hirozawa, MPH, Priscilla Lee Chu, MPH, Sharon Pipkin, MPH,
Tara Schubert, MS, Annie Vu, MPH
The AIDS Surveillance Report is accessible via internet: www.sfdph.org/PHP/HIVEpiSect.htm

AIDS Surveillance Summary

San Francisco (as of 06/30/07)

Cumulative cases ¹ :	27,234
Cumulative deaths:	18,420


California (as of 05/31/07)

Cumulative cases:	145,419
Cumulative deaths:	83,566

United States² (as of 12/31/05)

Cumulative cases:	988,376
Cumulative deaths:	550,394

San Francisco AIDS Incidence, Mortality, and Prevalence by Year, 1980-2007³



1. Includes SF residents diagnosed in SF and SF residents diagnosed in other jurisdictions. Excludes persons diagnosed in SF who resided in other jurisdictions at the time of their AIDS diagnosis.
2. The US numbers do not represent actual cases or deaths reported. Rather, these numbers are estimated and adjusted for reporting delays.
3. Reporting for recent year is incomplete. See Table 9 for actual numbers per year.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report

Summary of San Francisco Residents with AIDS* Reported as of 06/30/2007

Table 1. Adult/Adolescent AIDS Cases (>12 years) by Transmission Category, San Francisco, 1980-2007

Transmission Category	No.	(%)
Gay or bisexual male	20159	(74.1)
Heterosexual male injection drug user	1369	(5.0)
Heterosexual female injection drug user	664	(2.4)
Gay or bisexual male injection drug user	3831	(14.1)
Lesbian or bisexual injection drug user	43	(0.2)
Transgender (1)	360	(1.3)
Hemophiliac	16	(0.1)
Heterosexual contact male (2)	106	(0.4)
Heterosexual contact female (2)	266	(1.0)
Transfusion recipient	143	(0.5)
Risk not reported/Other (3)	239	(0.9)
Total	27196	(100)

Table 2. AIDS Cases by Gender and Year of Diagnosis, San Francisco, 1980-2007

Gender	Year of Diagnosis					
	< 1997 No. (%)	1997 No. (%)	1998 No. (%)	1999 No. (%)	2000 No. (%)	2001 No. (%)
Male	20898 (96.0)	737 (91.4)	625 (90.3)	513 (88.6)	479 (86.8)	443 (88.4)
Female	676 (3.1)	49 (6.1)	50 (7.2)	48 (8.3)	52 (9.4)	44 (8.8)
Transgender (1)	193 (0.9)	20 (2.5)	17 (2.5)	18 (3.1)	21 (3.8)	14 (2.8)
Total	21767 (100)	806 (100)	692 (100)	579 (100)	552 (100)	501 (100)

Gender	Year of Diagnosis					
	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)
Male	434 (89.1)	471 (88.5)	404 (88.4)	380 (89.0)	321 (89.4)	70 (93.3)
Female	35 (7.2)	38 (7.1)	39 (8.5)	34 (8.0)	29 (8.1)	5 (6.7)
Transgender (1)	18 (3.7)	23 (4.3)	14 (3.1)	13 (3.0)	9 (2.5)	0 (0.0)
Total	487 (100)	532 (100)	457 (100)	427 (100)	359 (100)	75 (100)

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Transgender information was collected since September 1996. Data prior to this are incomplete.

(2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.

(3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report

Summary of San Francisco Residents with AIDS* Reported as of 06/30/2007

Table 3. AIDS Cases by Transmission Category and Race/Ethnicity, San Francisco, 1980-2007

Transmission Category (1)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)
Adult/Adolescent					
Gay or bisexual male	15732 (80.4)	1514 (43.8)	2329 (73.5)	650 (76.6)	65 (45.8)
Injection drug user (IDU)	714 (3.6)	1026 (29.7)	231 (7.3)	43 (5.1)	21 (14.8)
Gay or bisexual male IDU	2844 (14.5)	645 (18.7)	416 (13.1)	65 (7.7)	49 (34.5)
Lesbian or bisexual IDU	21 (0.1)	14 (0.4)	5 (0.2)	2 (0.2)	1 (0.7)
Hemophiliac	8 (0.0)	2 (0.1)	5 (0.2)	1 (0.1)	0 (0.0)
Heterosexual (2)	111 (0.6)	146 (4.2)	82 (2.6)	31 (3.7)	3 (2.1)
Transfusion recipient	69 (0.4)	26 (0.8)	26 (0.8)	22 (2.6)	0 (0.0)
Risk not reported/Other (3)	70 (0.4)	69 (2.0)	66 (2.1)	30 (3.5)	2 (1.4)
Pediatric (0-12 years) (4)					
	8 (0.0)	13 (0.4)	10 (0.3)	5 (0.6)	1 (0.7)
Total	19577 (100)	3455 (100)	3170 (100)	849 (100)	142 (100)

Table 4. AIDS Cases by Transmission Category and Asian/Pacific Islander Ethnicity, San Francisco, 1980-2007

Transmission Category (1)	Chinese No. (%)	Japanese No. (%)	Filipino No. (%)	Southeast Asian No. (%)	Korean No. (%)	Pacific Islander No. (%)
Adult/Adolescent						
Gay or bisexual male	152 (79.2)	83 (83.8)	219 (80.5)	57 (71.3)	9 (75.0)	43 (65.2)
Injection drug user (IDU)	8 (4.2)	0 (0.0)	9 (3.3)	3 (3.8)	2 (16.7)	9 (13.6)
Gay or bisexual male IDU	8 (4.2)	12 (12.1)	19 (7.0)	7 (8.8)	1 (8.3)	7 (10.6)
Lesbian or bisexual IDU	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (3.0)
Hemophiliac	0 (0.0)	1 (1.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (2)	4 (2.1)	1 (1.0)	12 (4.4)	5 (6.3)	0 (0.0)	4 (6.1)
Transfusion recipient	11 (5.7)	1 (1.0)	7 (2.6)	3 (3.8)	0 (0.0)	0 (0.0)
Risk not reported/Other (3)	7 (3.6)	1 (1.0)	5 (1.8)	5 (6.3)	0 (0.0)	1 (1.5)
Pediatric (0-12 years) (4)						
	2 (1.0)	0 (0.0)	1 (0.4)	0 (0.0)	0 (0.0)	0 (0.0)
Total	192 (100)	99 (100)	272 (100)	80 (100)	12 (100)	66 (100)

* Residents of San Francisco at time of initial AIDS diagnosis.

- (1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.
- (4) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report

Summary of San Francisco Residents with AIDS* Reported as of 06/30/2007

Table 5. AIDS Cases by Transmission Category and Year of Diagnosis, San Francisco, 1980-2007

Transmission Category (1)	Year of Diagnosis					
	< 1997 No. (%)	1997 No. (%)	1998 No. (%)	1999 No. (%)	2000 No. (%)	2001 No. (%)
Adult/Adolescent						
Gay or bisexual male	16841 (77.4)	553 (68.6)	441 (63.7)	360 (62.2)	335 (60.7)	307 (61.3)
Injection drug user (IDU)	1296 (6.0)	106 (13.2)	91 (13.2)	81 (14.0)	88 (15.9)	68 (13.6)
Gay or bisexual male IDU	3112 (14.3)	115 (14.3)	136 (19.7)	103 (17.8)	98 (17.8)	83 (16.6)
Lesbian or bisexual IDU	29 (0.1)	2 (0.2)	2 (0.3)	1 (0.2)	3 (0.5)	3 (0.6)
Hemophiliac	15 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (2)	207 (1.0)	16 (2.0)	13 (1.9)	18 (3.1)	18 (3.3)	17 (3.4)
Transfusion recipient	138 (0.6)	2 (0.2)	0 (0.0)	1 (0.2)	0 (0.0)	1 (0.2)
Risk not reported/Other (3)	96 (0.4)	11 (1.4)	9 (1.3)	14 (2.4)	9 (1.6)	22 (4.4)
Pediatric (0-12 years) (4)	33 (0.2)	1 (0.1)	0 (0.0)	1 (0.2)	1 (0.2)	0 (0.0)
Total	21767 (100)	806 (100)	692 (100)	579 (100)	552 (100)	501 (100)

Transmission Category (1)	Year of Diagnosis					
	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)
Adult/Adolescent						
Gay or bisexual male	306 (62.8)	345 (64.8)	298 (65.2)	259 (60.7)	224 (62.4)	45 (60.0)
Injection drug user (IDU)	66 (13.6)	79 (14.8)	55 (12.0)	56 (13.1)	44 (12.3)	9 (12.0)
Gay or bisexual male IDU	89 (18.3)	74 (13.9)	76 (16.6)	73 (17.1)	59 (16.4)	10 (13.3)
Lesbian or bisexual IDU	0 (0.0)	2 (0.4)	0 (0.0)	0 (0.0)	1 (0.3)	0 (0.0)
Hemophiliac	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.3)	0 (0.0)
Heterosexual (2)	14 (2.9)	19 (3.6)	16 (3.5)	16 (3.7)	16 (4.5)	4 (5.3)
Transfusion recipient	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported/Other (3)	10 (2.1)	12 (2.3)	12 (2.6)	23 (5.4)	14 (3.9)	7 (9.3)
Pediatric (0-12 years) (4)	1 (0.2)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Total	487 (100)	532 (100)	457 (100)	427 (100)	359 (100)	75 (100)

* Residents of San Francisco at time of initial AIDS diagnosis.

- (1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.
- (4) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report

Summary of San Francisco Residents with AIDS* Reported as of 06/30/2007

Table 6. AIDS Cases by Gender, Age Group and Race/Ethnicity, San Francisco, 1980-2007

Male Age at Diagnosis (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	4 (0.0)	4 (0.1)	5 (0.2)	4 (0.5)	1 (0.8)	19 (0.1)
13 - 19	9 (0.0)	0 (0.0)	14 (0.5)	1 (0.1)	2 (1.6)	26 (0.1)
20 - 24	298 (1.6)	66 (2.3)	94 (3.2)	22 (2.9)	5 (4.0)	486 (1.9)
25 - 29	1625 (8.5)	247 (8.7)	414 (14.2)	85 (11.2)	23 (18.4)	2397 (9.3)
30 - 39	8666 (45.3)	1163 (41.0)	1413 (48.5)	341 (45.0)	62 (49.6)	11659 (45.2)
40 - 49	6231 (32.6)	919 (32.4)	711 (24.4)	228 (30.1)	26 (20.8)	8124 (31.5)
50 - 59	1830 (9.6)	346 (12.2)	208 (7.1)	59 (7.8)	5 (4.0)	2454 (9.5)
60 +	449 (2.3)	89 (3.1)	54 (1.9)	17 (2.2)	1 (0.8)	610 (2.4)
Male subtotal	19112 (100)	2834 (100)	2913 (100)	757 (100)	125 (100)	25775 (100)
Female Age at Diagnosis (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	4 (1.1)	9 (1.8)	5 (3.4)	1 (1.6)	0 (0.0)	19 (1.7)
13 - 19	1 (0.3)	1 (0.2)	2 (1.3)	0 (0.0)	0 (0.0)	4 (0.4)
20 - 24	14 (3.9)	9 (1.8)	10 (6.7)	4 (6.3)	1 (7.7)	38 (3.5)
25 - 29	40 (11.1)	42 (8.3)	23 (15.4)	11 (17.2)	1 (7.7)	118 (10.7)
30 - 39	147 (40.7)	202 (39.8)	48 (32.2)	20 (31.3)	8 (61.5)	429 (39.0)
40 - 49	95 (26.3)	174 (34.3)	39 (26.2)	19 (29.7)	3 (23.1)	330 (30.0)
50 - 59	31 (8.6)	47 (9.3)	13 (8.7)	6 (9.4)	0 (0.0)	97 (8.8)
60 +	29 (8.0)	23 (4.5)	9 (6.0)	3 (4.7)	0 (0.0)	64 (5.8)
Female subtotal	361 (100)	507 (100)	149 (100)	64 (100)	13 (100)	1099 (100)
Transgender (2) Age at Diagnosis (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander Native American No. (%)	Total (1) No. (%)	
13 - 29	22 (21.2)	25 (21.9)	29 (26.9)	9 (28.1)	85 (23.6)	
30 - 39	51 (49.0)	42 (36.8)	53 (49.1)	16 (50.0)	163 (45.3)	
40 +	31 (29.8)	47 (41.2)	26 (24.1)	7 (21.9)	112 (31.1)	
Transgender subtotal	104 (100)	114 (100)	108 (100)	32 (100)	360 (100)	

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Total includes persons with multiple or unknown race.

(2) Transgender information was collected since September 1996. Certain age or race/ethnic groups are combined for transgender cases because of small number.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report

Summary of San Francisco Residents with AIDS* Reported as of 06/30/2007

Table 7. AIDS Cases by Race/Ethnicity and Year of Diagnosis, San Francisco, 1980-2007

Race/Ethnicity	Year of Diagnosis					
	< 1997 No. (%)	1997 No. (%)	1998 No. (%)	1999 No. (%)	2000 No. (%)	2001 No. (%)
White	16369 (75.2)	515 (63.9)	443 (64.0)	339 (58.5)	319 (57.8)	291 (58.1)
African American	2434 (11.2)	146 (18.1)	140 (20.2)	104 (18.0)	112 (20.3)	100 (20.0)
Latino	2271 (10.4)	110 (13.6)	77 (11.1)	104 (18.0)	88 (15.9)	70 (14.0)
Asian/Pacific Islander	578 (2.7)	30 (3.7)	28 (4.0)	25 (4.3)	27 (4.9)	31 (6.2)
Native American	102 (0.5)	4 (0.5)	2 (0.3)	5 (0.9)	4 (0.7)	5 (1.0)
Total (1)	21767 (100)	806 (100)	692 (100)	579 (100)	552 (100)	501 (100)

Race/Ethnicity	Year of Diagnosis					
	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)
White	290 (59.5)	271 (50.9)	255 (55.8)	246 (57.6)	195 (54.3)	44 (58.7)
African American	86 (17.7)	102 (19.2)	75 (16.4)	76 (17.8)	69 (19.2)	11 (14.7)
Latino	74 (15.2)	118 (22.2)	100 (21.9)	78 (18.3)	65 (18.1)	15 (20.0)
Asian/Pacific Islander	29 (6.0)	31 (5.8)	19 (4.2)	24 (5.6)	22 (6.1)	5 (6.7)
Native American	3 (0.6)	6 (1.1)	4 (0.9)	2 (0.5)	5 (1.4)	0 (0.0)
Total (1)	487 (100)	532 (100)	457 (100)	427 (100)	359 (100)	75 (100)

Table 8. AIDS Cases and Cumulative Rates per 100,000 by Race/Ethnicity and Gender, San Francisco

Race/Ethnicity	Male		Female		Total (2)	
	No.	(Rate)	No.	(Rate)	No.	(Rate)
White	19112	(11044.0)	361	(220.0)	19473	(5776.3)
African American	2834	(7603.0)	507	(1297.7)	3341	(4376.3)
Latino	2913	(5676.9)	149	(301.6)	3062	(3040.2)
Asian/Pacific Islander	757	(766.7)	64	(59.8)	821	(399.2)
Native American	125	(9097.5)	13	(1030.9)	138	(5237.2)
Total (1)	25775	(7110.4)	1099	(304.0)	26874	(3712.1)

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Total includes persons with multiple or unknown race.

(2) Transgender cases were excluded because population size for transgender can not be determined for rates calculation.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report

Summary of San Francisco Residents with AIDS* Reported as of 06/30/2007

Table 9. AIDS Incidence, Mortality, and Prevalence by Year, San Francisco, 1980-2007

Year	Number of Cases Reported per Year	Number of Cases Diagnosed per Year (1)	Number of Deaths Occurred per Year (1)	Number of Persons Living with AIDS (1)
1980	0	3	0	3
1981	21	26	8	21
1982	75	99	32	88
1983	197	274	111	251
1984	451	557	273	535
1985	673	859	532	862
1986	981	1236	807	1291
1987	1287	1629	877	2043
1988	1408	1762	1038	2767
1989	1585	2161	1275	3653
1990	1687	2048	1364	4337
1991	1685	2283	1504	5116
1992	1639	2326	1641	5801
1993	4272	2074	1599	6276
1994	1916	1786	1592	6470
1995	1635	1561	1483	6548
1996	1246	1083	987	6644
1997	1061	806	422	7028
1998	795	692	401	7319
1999	726	579	355	7543
2000	628	552	346	7749
2001	498	501	322	7928
2002	441	487	324	8091
2003	533	532	301	8322
2004	558	457	304	8475
2005	516	427	248	8654
2006	440	359	206	8807
2007	280	75	68	8814
Total	27234	27234	18420	

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Data in recent years is incomplete due to delay in cases/deaths reporting.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report

Summary of San Francisco Residents with AIDS* Reported as of 06/30/2007

Table 10. Cases by Initial AIDS-Defining Condition, San Francisco, 1980-2007

Initial AIDS-Defining Condition	Def. (1) No.	Pres. (2) No.	Total No. (%)
Bacterial infections, recurrent, <13 years [HIV+(3)]	8	N/A(4)	8 (0.0)
Candidiasis of bronchi, trachea, or lungs	24	N/A	24 (0.1)
Candidiasis of esophagus	294	220	514 (1.9)
Cervical cancer, invasive [HIV+]	5	N/A	5 (0.0)
Coccidioidomycosis, disseminated or extrapulmonary [HIV+]	7	N/A	7 (0.0)
Cryptococcosis, extrapulmonary	372	N/A	372 (1.4)
Cryptosporidiosis, intestinal (>1 mo. duration)	331	N/A	331 (1.2)
Cytomegalovirus (except liver, spleen, lymph nodes), >1 month of age	202	N/A	202 (0.7)
CMV retinitis with loss of vision [HIV+]	0	113	113 (0.4)
HIV encephalopathy [HIV+]	382	N/A	382 (1.4)
Herpes simplex: chronic (>1 mo.), bronchitis, pneumonitis, esophagitis	96	N/A	96 (0.4)
Histoplasmosis, disseminated or extrapulmonary [HIV+]	25	N/A	25 (0.1)
Isosporiasis, intestinal (>1 mo. duration) [HIV+]	21	N/A	21 (0.1)
Kaposi's sarcoma (5)	2536	281	2817 (10.3)
Lymphoid interstitial pneumonia/pulmonary lymphoid hyperplasia, <13 years	5	0	5 (0.0)
Lymphoma, Burkitt's (non-Hodgkin's) [HIV+]	103	N/A	103 (0.4)
Lymphoma, immunoblastic (non-Hodgkin's) [HIV+]	415	N/A	415 (1.5)
Lymphoma, primary in brain (5)	40	N/A	40 (0.1)
Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary	254	3	257 (0.9)
Mycobacterium tuberculosis, pulmonary [HIV+]	195	13	208 (0.8)
Mycobacterium tuberculosis, disseminated or extrapulmonary [HIV+]	142	5	147 (0.5)
Mycobacterium other species, disseminated or extrapulmonary [HIV+]	16	11	27 (0.1)
Pneumocystis carinii pneumonia	5068	589	5657 (20.8)
Pneumonia, recurrent [HIV+]	219	24	243 (0.9)
Progressive multifocal leukoencephalopathy	50	N/A	50 (0.2)
Salmonella sepsis, recurrent [HIV+]	8	N/A	8 (0.0)
Toxoplasmosis of brain, >1 month of age	44	199	243 (0.9)
Wasting syndrome [HIV+]	675	N/A	675 (2.5)
CD4 T lymphocyte count <200 or percent <14 [HIV+]	14235	N/A	14235 (52.3)
Any AIDS indicator condition and HIV-negative and CD4 count <400 (6)	4	N/A	4 (0.0)
Total	25776	1458	27234 (100)

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Indicator conditions diagnosed definitively (e.g. culture or biopsy proven).

(2) Indicator conditions diagnosed presumptively in a person who has laboratory evidence of HIV infection.

(3) [HIV+]: Indicator conditions that require laboratory evidence of HIV infection.

(4) N/A: Conditions which require definitive diagnoses only.

(5) Laboratory evidence of HIV infection in persons > 60 years of age.

(6) In the absence of other causes of immunocompromise.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report

Summary of San Francisco Residents with AIDS* Reported as of 06/30/2007

Table 11. Cumulative AIDS Indicator Conditions among Persons with AIDS, San Francisco, 1980-2007

AIDS Indicator Condition (1)	Total No. (%)
Bacterial infections, recurrent, <13 years [HIV+(2)]	11 (0.0)
Candidiasis of bronchi, trachea, or lungs	135 (0.5)
Candidiasis of esophagus	2469 (9.1)
Cervical cancer, invasive [HIV+]	11 (0.0)
Coccidioidomycosis, disseminated or extrapulmonary [HIV+]	59 (0.2)
Cryptococcosis, extrapulmonary	1801 (6.6)
Cryptosporidiosis, intestinal (>1 mo. duration)	1363 (5.0)
Cytomegalovirus (except liver, spleen, lymph nodes), >1 month of age	2414 (8.9)
CMV retinitis with loss of vision [HIV+]	2472 (9.1)
HIV encephalopathy [HIV+]	2452 (9.0)
Herpes simplex: chronic (>1 mo.), bronchitis, pneumonitis, esophagitis	408 (1.5)
Histoplasmosis, disseminated or extrapulmonary [HIV+]	154 (0.6)
Isosporiasis, intestinal (>1 mo. duration) [HIV+]	67 (0.2)
Kaposi's sarcoma (3)	6630 (24.3)
Lymphoid interstitial pneumonia/pulmonary lymphoid hyperplasia, <13 years	5 (0.0)
Lymphoma, Burkitt's (non-Hodgkin's) [HIV+]	532 (2.0)
Lymphoma, immunoblastic (non-Hodgkin's) [HIV+]	1113 (4.1)
Lymphoma, primary in brain (3)	385 (1.4)
Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary	5035 (18.5)
Mycobacterium tuberculosis, pulmonary [HIV+]	629 (2.3)
Mycobacterium tuberculosis, disseminated or extrapulmonary [HIV+]	469 (1.7)
Mycobacterium other species, disseminated or extrapulmonary [HIV+]	326 (1.2)
Pneumocystis carinii pneumonia	10828 (39.8)
Pneumonia, recurrent [HIV+]	982 (3.6)
Progressive multifocal leukoencephalopathy	300 (1.1)
Salmonella sepsis, recurrent [HIV+]	57 (0.2)
Toxoplasmosis of brain, >1 month of age	1150 (4.2)
Wasting syndrome [HIV+]	4291 (15.8)

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Cases may have more than one condition.

(2) [HIV+]: Indicator conditions that require laboratory evidence of HIV infection.

(3) Laboratory evidence of HIV infection in persons > 60 years of age.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report

Summary of San Francisco Residents with AIDS* Reported as of 06/30/2007

Table 12. Living Adult/Adolescent AIDS Cases (>12 years) by Transmission Category, San Francisco

Transmission Category	No.	(%)
Gay or bisexual male	6311	(71.7)
Heterosexual male injection drug user	481	(5.5)
Heterosexual female injection drug user	283	(3.2)
Gay or bisexual male injection drug user	1159	(13.2)
Lesbian or bisexual injection drug user	20	(0.2)
Transgender (1)	187	(2.1)
Hemophiliac	5	(0.1)
Heterosexual contact male (2)	60	(0.7)
Heterosexual contact female (2)	146	(1.7)
Transfusion recipient	21	(0.2)
Risk not reported/Other (3)	127	(1.4)
Total	8800	(100)

Table 13. Living AIDS Cases by Transmission Category and Race/Ethnicity, San Francisco

Transmission Category	White		African American		Latino		Asian/Pacific Islander		Native American	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Adult/Adolescent										
Gay or bisexual male	4536	(79.1)	508	(41.1)	1011	(74.1)	301	(74.3)	23	(44.2)
Injection drug user (IDU)	276	(4.8)	375	(30.4)	84	(6.2)	19	(4.7)	10	(19.2)
Gay or bisexual male IDU	813	(14.2)	224	(18.1)	168	(12.3)	34	(8.4)	15	(28.8)
Lesbian or bisexual IDU	10	(0.2)	8	(0.6)	1	(0.1)	1	(0.2)	0	(0.0)
Hemophiliac	3	(0.1)	2	(0.2)	0	(0.0)	0	(0.0)	0	(0.0)
Heterosexual (2)	53	(0.9)	76	(6.2)	55	(4.0)	20	(4.9)	2	(3.8)
Transfusion recipient	7	(0.1)	3	(0.2)	5	(0.4)	6	(1.5)	0	(0.0)
Risk not reported/Other (3)	34	(0.6)	35	(2.8)	35	(2.6)	21	(5.2)	1	(1.9)
Pediatric (0-12 years) (4)	0	(0.0)	4	(0.3)	5	(0.4)	3	(0.7)	1	(1.9)
Total	5732	(100)	1235	(100)	1364	(100)	405	(100)	52	(100)

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Transgender information was collected since September 1996. Data prior to this are incomplete.

(2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.

(3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.

(4) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report

Summary of San Francisco Residents with AIDS* Reported as of 06/30/2007

Table 14. Living AIDS Cases by Gender, Race/Ethnicity and Age as of 06/30/2007, San Francisco

Male Current Age (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	0 (0.0)	0 (0.0)	1 (0.1)	0 (0.0)	0 (0.0)	2 (0.0)
13 - 19	0 (0.0)	0 (0.0)	0 (0.0)	2 (0.6)	1 (2.2)	3 (0.0)
20 - 24	5 (0.1)	3 (0.3)	9 (0.7)	2 (0.6)	1 (2.2)	20 (0.2)
25 - 29	31 (0.6)	7 (0.7)	33 (2.7)	6 (1.7)	3 (6.7)	81 (1.0)
30 - 39	562 (10.2)	109 (11.4)	255 (20.8)	76 (21.7)	9 (20.0)	1013 (12.5)
40 - 49	2310 (41.7)	389 (40.8)	575 (47.0)	143 (40.7)	23 (51.1)	3452 (42.4)
50 - 59	1954 (35.3)	331 (34.7)	257 (21.0)	95 (27.1)	8 (17.8)	2649 (32.6)
60 +	674 (12.2)	115 (12.1)	94 (7.7)	27 (7.7)	0 (0.0)	912 (11.2)
Male subtotal	5536 (100)	954 (100)	1224 (100)	351 (100)	45 (100)	8132 (100)
Female Current Age (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	0 (0.0)	1 (0.5)	1 (1.2)	0 (0.0)	0 (0.0)	2 (0.4)
13 - 19	0 (0.0)	2 (0.9)	2 (2.5)	1 (2.9)	0 (0.0)	5 (1.0)
20 - 24	0 (0.0)	1 (0.5)	2 (2.5)	0 (0.0)	0 (0.0)	3 (0.6)
25 - 29	1 (0.7)	1 (0.5)	5 (6.2)	6 (17.1)	1 (20.0)	14 (2.8)
30 - 39	28 (18.8)	44 (19.8)	16 (19.8)	8 (22.9)	0 (0.0)	99 (20.0)
40 - 49	64 (43.0)	77 (34.7)	30 (37.0)	10 (28.6)	3 (60.0)	184 (37.2)
50 - 59	47 (31.5)	75 (33.8)	17 (21.0)	8 (22.9)	1 (20.0)	148 (29.9)
60 +	9 (6.0)	21 (9.5)	8 (9.9)	2 (5.7)	0 (0.0)	40 (8.1)
Female subtotal	149 (100)	222 (100)	81 (100)	35 (100)	5 (100)	495 (100)
Transgender (2) Current Age (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander Native American No. (%)	Total (1) No. (%)	
13 - 39	15 (31.9)	10 (16.9)	22 (37.3)	6 (28.6)	53 (28.3)	
40 +	32 (68.1)	49 (83.1)	37 (62.7)	15 (71.4)	134 (71.7)	
Transgender subtotal	47 (100)	59 (100)	59 (100)	21 (100)	187 (100)	

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Total includes persons with multiple or unknown race.

(2) Transgender information was collected since September 1996. Certain age or race/ethnic groups are combined for transgender cases because of small number.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report

Summary of San Francisco Residents with AIDS* Reported as of 06/30/2007

Table 15. Living AIDS Cases by Initial AIDS-Defining Condition, San Francisco

Initial AIDS-Defining Condition	Def. (1) No.	Pres. (2) No.	Total No. (%)
Bacterial infections, recurrent, <13 years [HIV+(3)]	7	N/A(4)	7 (0.1)
Candidiasis of bronchi, trachea, or lungs	5	N/A	5 (0.1)
Candidiasis of esophagus	45	52	97 (1.1)
Cervical cancer, invasive [HIV+]	0	N/A	0 (0.0)
Coccidioidomycosis, disseminated or extrapulmonary [HIV+]	1	N/A	1 (0.0)
Cryptococcosis, extrapulmonary	45	N/A	45 (0.5)
Cryptosporidiosis, intestinal (>1 mo. duration)	108	N/A	108 (1.2)
Cytomegalovirus (except liver, spleen, lymph nodes), >1 month of age	24	N/A	24 (0.3)
CMV retinitis with loss of vision [HIV+]	0	22	22 (0.2)
HIV encephalopathy [HIV+]	59	N/A	59 (0.7)
Herpes simplex: chronic (>1 mo.), bronchitis, pneumonitis, esophagitis	14	N/A	14 (0.2)
Histoplasmosis, disseminated or extrapulmonary [HIV+]	8	N/A	8 (0.1)
Isosporiasis, intestinal (>1 mo. duration) [HIV+]	7	N/A	7 (0.1)
Kaposi's sarcoma (5)	323	61	384 (4.4)
Lymphoid interstitial pneumonia/pulmonary lymphoid hyperplasia, <13 years	3	0	3 (0.0)
Lymphoma, Burkitt's (non-Hodgkin's) [HIV+]	21	N/A	21 (0.2)
Lymphoma, immunoblastic (non-Hodgkin's) [HIV+]	47	N/A	47 (0.5)
Lymphoma, primary in brain (5)	2	N/A	2 (0.0)
Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary	17	0	17 (0.2)
Mycobacterium tuberculosis, pulmonary [HIV+]	78	5	83 (0.9)
Mycobacterium tuberculosis, disseminated or extrapulmonary [HIV+]	37	2	39 (0.4)
Mycobacterium other species, disseminated or extrapulmonary [HIV+]	2	2	4 (0.0)
Pneumocystis carinii pneumonia	407	138	545 (6.2)
Pneumonia, recurrent [HIV+]	83	8	91 (1.0)
Progressive multifocal leukoencephalopathy	0	N/A	0 (0.0)
Salmonella sepsis, recurrent [HIV+]	1	N/A	1 (0.0)
Toxoplasmosis of brain, >1 month of age	9	22	31 (0.4)
Wasting syndrome [HIV+]	137	N/A	137 (1.6)
CD4 T lymphocyte count <200 or percent <14 [HIV+]	7011	N/A	7011 (79.5)
Any AIDS indicator condition and HIV-negative and CD4 count <400 (6)	1	N/A	1 (0.0)
Total	8502	312	8814 (100)

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Indicator conditions diagnosed definitively (e.g. culture or biopsy proven).

(2) Indicator conditions diagnosed presumptively in a person who has laboratory evidence of HIV infection.

(3) [HIV+]: Indicator conditions that require laboratory evidence of HIV infection.

(4) N/A: Conditions which require definitive diagnoses only.

(5) Laboratory evidence of HIV infection in persons > 60 years of age.

(6) In the absence of other causes of immunocompromise.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report

Summary of San Francisco Residents with AIDS* Reported as of 06/30/2007

Table 16. Cumulative AIDS Indicator Conditions among Persons Living with AIDS, San Francisco

AIDS Indicator Condition (1)	Total No. (%)
Bacterial infections, recurrent, <13 years [HIV+(2)]	10 (0.1)
Candidiasis of bronchi, trachea, or lungs	23 (0.3)
Candidiasis of esophagus	424 (4.8)
Cervical cancer, invasive [HIV+]	1 (0.0)
Coccidioidomycosis, disseminated or extrapulmonary [HIV+]	12 (0.1)
Cryptococcosis, extrapulmonary	254 (2.9)
Cryptosporidiosis, intestinal (>1 mo. duration)	302 (3.4)
Cytomegalovirus (except liver, spleen, lymph nodes), >1 month of age	199 (2.3)
CMV retinitis with loss of vision [HIV+]	201 (2.3)
HIV encephalopathy [HIV+]	199 (2.3)
Herpes simplex: chronic (>1 mo.), bronchitis, pneumonitis, esophagitis	67 (0.8)
Histoplasmosis, disseminated or extrapulmonary [HIV+]	24 (0.3)
Isosporiasis, intestinal (>1 mo. duration) [HIV+]	15 (0.2)
Kaposi's sarcoma (3)	876 (9.9)
Lymphoid interstitial pneumonia/pulmonary lymphoid hyperplasia, <13 years	2 (0.0)
Lymphoma, Burkitt's (non-Hodgkin's) [HIV+]	62 (0.7)
Lymphoma, immunoblastic (non-Hodgkin's) [HIV+]	118 (1.3)
Lymphoma, primary in brain (3)	13 (0.1)
Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary	303 (3.4)
Mycobacterium tuberculosis, pulmonary [HIV+]	190 (2.2)
Mycobacterium tuberculosis, disseminated or extrapulmonary [HIV+]	101 (1.1)
Mycobacterium other species, disseminated or extrapulmonary [HIV+]	42 (0.5)
Pneumocystis carinii pneumonia	1380 (15.7)
Pneumonia, recurrent [HIV+]	251 (2.8)
Progressive multifocal leukoencephalopathy	13 (0.1)
Salmonella sepsis, recurrent [HIV+]	3 (0.0)
Toxoplasmosis of brain, >1 month of age	81 (0.9)
Wasting syndrome [HIV+]	655 (7.4)

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Cases may have more than one condition.

(2) [HIV+]: Indicator conditions that require laboratory evidence of HIV infection.

(3) Laboratory evidence of HIV infection in persons > 60 years of age.